

TAXONOMIC INDEX

Copeia 2012, Nos. 1–4

NEW NAMES

Adontosternarchus duartei 535–540
Amolops indoburmanensis 58, 60, 62–66, 68, 70–74
Astyanacinus yariquies 501–506
Apteronotus lindalvae 283–292
Bathyaethiops atercrinis 478–483
Careproctus kamikawai 257–265
lycopersicus 257–265
Cathorops wayuu 79, 81–84, 88, 90–94
Congoglanis howesi 626–630
Ctenocheiroidon pristis 243–250
Derilissus lombardii 251–253, 255–256
Guibemantis tasifotsy 648–662
Hisonotus bockmanni 266–275
Hypostomus careopinnatus 494–500
Loricaria coximensis 49–56

Nannoplectostomus 637–647
eleonora 637–647
Pareiorhaphis proskynita 519–524
Peckoltia feldbergae 547–553
Plesiopoma 6–11
curvidens 6–11
Potamorhaphis labiatus 663–668
Rypticus carpenteri 24–30, 32–35
Scinax imbegue 554–569
tymbamirim 554–569
Scoloplax baileyi 670–677
Tricheilostoma jani 39–45, 47
Trichomycterus minus 631–636
Trinectes hubbsbollinger 541–546

NEW AND ESTABLISHED NAMES

Abudefduf sp. 217
Acanthochromis polyacanthus 217
Acanthophis barnetti 578
bottomi 578
crotalusei 578
cumingi 578
macgregori 578
wellsi 578, 580
woolfi 578
Acari 121–129
Acarina 205, 207
Acentrogobius suluensis 533
Acestrorhynchinae 618
Acheta domesticus 145
Achiridae 541–546
Acinocheriodon 243
Acipenser baeri 390
Acris crepitans 191–196, 678
c. blanchardi 154
Acropora 217–218
Actinopterygii 165
Acyrtops artius 256
beryllinus 256
rubiginosus 256
Adamsia 204
Adenomera horrida 205
Adinia xenica 451, 456
Adolfus vauereselli 580
Adontosternarchus 535–540
balaenops 536–537, 539
cf. sachsi 540
clarkae 535–537, 539

devenanzii 535, 537, 539–540
duartei 535–540
nebulosus 537, 539–540
sachsi 537, 539–540
Adrianichthyidae 776
Aechmea nudicaulis 561, 566
Aframomum 655
Agalychnis callidryas 198
Agama neumanni 579
Agamidae 416
Agkistrodon piscivorus 358
Aipysurus pooleorum 581
Alcantarea glaziouana 683–688
Alestidae 478–483
Algae 685–686
Alligator mississippiensis 359, 419–423
Allophrynidae 378
Allophryne ruthveni 378
Alona azorica 757, 759
Alopias superciliosus 765
vulpinus 765
Alsophis vudii 582
Amazonspither dalmata 243
Amblyglyphidodon curacao 217
Amblyopinae 527–534
Amblyraja radiata 786
Ambystoma 570–577, 786
annulatum 570–577
barbouri 573
californiense 573, 575
jeffersonianum 573, 575
laterale 573
macroactylum 331–340
m. croceum 331, 336

- m. macrodactylum* 573
maculatum 112, 312, 331, 336, 575
mavortium 331, 334
opacum 573, 575
talpoideum 112, 573, 575
texanum 332, 573, 575
tigrinum 573, 575
t. tigrinum 332
Amia 395
 calva 604–608
Amiidae 603, 606–607
Amolops 57, 59, 61, 63
 afghanus 57–76
 akhaorum 57, 65, 69, 72
 aniquiaoensis 57, 65, 69, 72
 archotaphus 57, 65, 69, 72
 assamanensis 69, 72
 assamensis 73
 bellulus 57, 60, 62, 65, 69, 72
 caelumnocis 65, 69, 72
 chakrataensis 65, 69, 72
 chunganensis 59–62, 65, 69, 72
 compotrix 65, 69, 72
 cremnobatus 60–62, 65, 69, 72
 cuciae 65, 69, 72
 daiyunensis 60, 62, 65, 69, 72
 daorum 65, 69, 72
 formosus 65, 69, 72
 gerbillus 57, 65, 69, 72
 granulosus 65, 69, 72
 hainanensis 60, 62, 65, 69, 72
 himalayanus 65, 69, 72
 hongkongensis 60, 62, 65, 69, 72
 indoburmanensis 58, 60, 62–66, 68, 70–74
 iriodes 65, 69, 72
 jinjiangensis 60–62, 65, 69, 72
 kantingensis 59–62, 65, 69, 72
 kaulbacki 57, 65, 69, 72
 kohimaensis 57, 65, 69, 72
 larutensis 65, 69, 72
 liangshanensis 60–62, 65, 69, 72
 lifanensis 60–62, 65, 69, 72
 loloensis 60–62, 65, 69, 72
 longimanus 57, 65, 69, 72
 mantzorum 57, 60–62, 65, 69, 72
 marmoratus 57–76
 medogensis 57, 65, 69, 72
 mengyangensis 65, 69, 72
 minutus 65, 69, 72
 monticola 65, 69, 72
 nasicus 65, 69, 72
 nidorbellus 57, 65, 69, 72
 panhai 57–76
 ricketti 60, 62, 65, 69, 72
 spinapectoralis 60, 62, 65, 69, 72
 splendissimus 65, 69, 72
 tormotus 65, 69, 72
 torrentis 60–62, 65, 69, 72
 viridimaculatus 57, 61–62, 65, 69, 72
 vitrea 65, 69, 72
 wuyiensis 65, 69, 72
Amphiesma groundwateri 581
Amphiglossus polleni 581
Amphiliidae 626
Amphipoda 121–129
Anabaena 686
Anaxyrus 168
 boreas 334
 cognatus 321–330
 fowleri 307–320
 woodhousii 321–330
Ancistomus 552
Ancistrini 547–553, 642
Ancistrus 524
 jataiensis 499, 524
 parecis 524
 reisi 499, 524, 645
 snethlageae 552
 sp. 551
 tombador 498–499, 524
 verecundus 524
Anguilliformes 484–493
Anolis 584–585, 770
 bicaorum 578
 bransfordii 581
 breedlovei 581
 (*Chamaeleolis*) 584
 (*Chamaelinorops*) 584
 chamaeleonides 584
 gorgonae 584
 greyi 581
 (*Norops*) 584
 proboscis 584
 sagrei 585
Anomaloglossus beebei 688
Anomalopus mackayi 581
Anomopoda 757–759
Anoplolepis gracilipes 127
Anostraca 757–759
Anotheca spinosa 688
Antaresia 578
Anthurium 723, 725
 sp. 723
Anura 106–114, 124, 168, 507–518, 722–731
Apalone 304–305
 mutica 305
 spinifera 305, 772
Aphanotorulus 494
Aphyocharacinae 618, 620
Aphyocharax 611
Aplastodiscus perviridis 514
Apogon
 frailis 386
 leptacanthus 386
 notatus 386
Aprasia 580
 haroldi 580
Apristurus microps 765
Apterodontidae 283–292, 535–540
Apterodontus 283–292
 albifrons 283–292
 brasiliensis 283–284, 286, 289
 camposdapazi 283–284, 286, 288
 caudimaculosus 283–284, 286, 288, 290
 cf. *albifrons* 290
 cf. *camposdapazi* 290
 cf. *eschmeyeri* 290
 cuchillejo 284, 286–287, 290

- cuchillo* 284, 286, 288, 290
ellisi 284, 286, 289
eschmeyer 283–284, 286, 288–290
galvisi 283–284, 286, 288
jurubidae 284, 286, 290
leptorhynchus 283–284, 286–287, 290
lindalvae 283–292
macrostomus 284, 286–287, 289–290
magdalenensis 284, 286–290
magoi 284, 286
mariae 284, 286, 289–290
milesi 283–284, 286, 288, 290
rostratus 283–284, 286, 290
sp. 291
spurrellii 284, 286–287
 Araceae 723
 Arachnida 121, 124–125, 205, 207
 Araneae 121–129, 205, 207
Arapaima 770
Arcella 685
Archocentrus octofasciatus 164
Archolaemus janeae 551
Arcos macrophthalmus 256
 Archosauria 772
Argyropelecus 395, 401
 sp. 395, 401, 405
 Ariidae 77–97
Aristida stricta 706
Aristostomias 395, 398–399, 405
 xenostoma 405
 Arius
 arenatus 94
 fissus 91, 94
 laticeps 77–97
 melanopus 95
 nigricans 77–97
 nuchalis 77–78, 81
 pleurops 91, 94
 punctulatus 77, 91
 spixii 78, 81, 87
 variolosus 77–97
Aronia melanocarpa 182
Artemia salina 343
Ascapheus 168
Aspidites melanocephalus 578
Aspidoscelis tigris 102
Astatoreochromis alluaudi 386
Astraptes 34
 fulgurator 34
Astroblepidae 631, 675
Astroblepus 631
Astronesthes 394–407
 ijimai 404–405
 indicus 405
 lucifer 395, 397–398, 403, 405
 niger 395, 397–398, 405
Astyanacinus 501–506
 moori 502
 moorii 501–502
 mulitdens 501–502
 n. sp. Bolivia 502
 n. sp. Peru 502
 n. sp. Purus 502
 yariquies 501–506
Astyanax 163, 501–502, 619
 atratoensis 505
 fasciatus 163
 filiferus 505
 leopoldi 501
 magdalenae 505
 microlepis 505
 orthodus 501–502, 504–505
 scabripinnis 523
 sp. 1 505
 superbus 501–502, 504
 villwocki 501
Atelopus oxyrhynchus 447
Atherinops affinis 341–350
 Atherinopsidae 166
 Atractus
 gaigeae 770
 lasallei 578
Auchenorrhyncha 124–126
Australoheros sp. 523
Awaous 293–300
 banana 294–297, 299
 grammepomus 297
 guamensis 293–300
 melanocephalus 297
 ocellaris 293–297, 299
 stamineus 293–300
Bachia 580
Baldellia ranunculoides 757
 Barbus
 cf. aloyi 481
 holoania 481
 jae 481
 kuiluensis 481
Baryancistrus 552
 xanthellus 551
Basiliscus plumifrons 465–471
Batagur baska 772
Bathophilus 395, 398–399, 405
 filifer 405
 pawnee 405
 sp. 405
 vaillanti 403, 405
Bathyaethiops 478–483
 atercrinis 478–483
 breuseghemi 478–479, 482
 caudomaculatus 478–479, 481, 483
 greeni 478–479, 481, 483
Bathyraja 34
Batrachochytrium dendrobatidis 150–159, 786
 Bdelloidea 685
 Belonidae 663–669
 Beloniformes 663–669, 776
Belonion 663
Betula pendula 182
 Bitis
 caudalis 735
 gabonica 735
 namaquensis 117
 schneideri 115–120, 732–737
 Blattodea 121, 125
Blommersia angolafa 648

Boa

- constrictor* 359, 473
- c. constrictor* 472–477
- c. imperator* 472–477
- c. occidentalis* 413, 472

Boidae 472–477**Boiga**

- andamanensis* 578
- irregularis* 169

Bokermannohyla

- aff. *circumdata* 523
- ahenea* 514
- nanuzae* 523

Bolitoglossa 171**Bolyeria** 580**Borostomias** 394–407

- elucens* 405
- panamensis* 402

Bothriopsis 770**Bothrops** 409, 413, 770

- alcatraz* 409
- jararaca* 409, 416
- insularis* 408–418

Brachionus 685**Brachyamblyopus** 527–534

- burmanicus* 530, 533
- intermedius* 530

Brachymeles

- lukbani* 581
- pathfinderi* 578

Brachypetersius 478–479

- altus* 478–483
- cadwaladeri* 479, 481
- gabonensis* 479, 481
- huloti* 479, 483
- notospilus* 479, 481, 483
- pseudonummifer* 479, 481, 483

Bramocharax 611**Branchiopoda** 757–759**Brasilia aequatoriali** 86**Brittanichthys** 22

- axelrodi* 619

Brogammerus

- reticulatus*
- r. patrickcouperi* 578
- r. stuartbigmorei* 578

Bromeliaceae 683, 685**Bryconamericus** sp. 523**Bryconinae** 618–619**Bryconops** 619**Buergeria japonica** 110, 112**Bufo** 168, 308

- americanus* 312
- boreas* 154
- bufo* 429, 447
- melanostictus* 112
- nebulifer* 106
- viridis* 429
- woodhousii* 312

Bufonidae 106–114, 378**Buhoma vauerocegae** 580**Bulbochaete** 514**Bullockia** 631, 634

- maldonadoi* 635

Cacophis churchilli 578†**Calamopleurus** 607

- africanus* 603–604, 607
- cylindricus* 607

Calanoida 757–759**Calathea** 723, 725

- sp. 723–724

Caledoniscincus orestes 580**Callagur borneoensis** 772**Callichthyidae** 675**Callinectes sapidus** 419–423**Callitris** 130**Calotes**

- emma* 582
- mariae* 582

Canis latrans 184**Caragobius urolepis** 533**Carcharinidae** 138, 766**Carcharhiniformes** 98–99**Carcharhinus**

- cerdale* 766
- imamus* 788
- perezii* 138

Carcharodon carcharias 766**Careproctus** 257–265

- cyclocephalus* 264
- kamikawai** 257–265
- lycopersicus** 257–265
- macrodiscus* 264
- ovigerus* 257–265

Caretta caretta 701**Carettochelys** 772**Carlia** 580

- amax* 578

Castor canadensis 179–190**Casuarina equisetifolia** 121**Cathorops** 77–97

- agassizii* 77, 79, 81–84, 88, 91–94
- aguadulce* 79, 83, 88, 93–94
- arenatus* 77–97
- belizensis* 79, 83, 88, 93–94
- cf. *spixii* 92
- dasycephalus* 77
- fuerthii* 80, 84, 89
- higuchii* 79, 83, 88, 93–94
- hypophthalmus* 80, 84, 89
- kailolae* 79, 83, 88, 93, 95
- laticeps* 91
- liropus* 80, 84, 89
- manglarensis* 80, 84, 89
- mapale* 77–97
- melanopus* 79, 83, 88, 93, 95
- multiradiatus* 80, 84, 89
- nuchalis* 77–97
- raredonae* 80, 84, 89
- sp. 87
- spixii* 77–97
- steindachneri* 80, 84, 89
- taylori* 80, 85, 89
- tuyra* 80, 85, 89
- wayuu** 79, 81–84, 88, 90–94

Catonotus 432–439**Celsiella** 729**Centrolenidae** 171, 722–731

- Centrophoridae 766
Centrophorus 766
 uyato 766
Centropyge potteri 777
Centrosymnus 766
Cephalopholis sp. 217
Ceratophora erdeleni 581
 Ceratopogonidae 634
Ceratoscopus maderensis 160
Ceriodaphnia sp. 757
 Cetorhinidae 766
Cetorhinus maximus 765
Chaenogobius 533
Chaetodon 211–212, 215, 219–220
 aureofasciatus 211–221
 auriga 211–221
 plebeius 211–221
 rainfordi 211–221
 sp. 213
 Chaetodontidae 211–221
Chamaeleo roperi 581
Channa 347
 Characidae 243–250, 478–483, 501–506, 609, 611, 618–619,
 621–622, 637
Characidium sp. 523
 Characiformes 501–506, 609, 619, 637
 Characinae 618–619
Charina bottae 103
Chasmistes muriei 167
Chauliodus 394–395, 398–399, 405, 583
 barbatus 397, 405
 eximius 582
 macouni 400, 405
 pammelas 404–405
 sloani 395, 397, 403, 405
Cheiridodus 494
Cheirodon 243, 245
 Cheirodontinae 243, 245, 248, 611, 618–622
 Cheirodontini 243–250
Chelmon 212
 rostratus 211–221
Chelodina canni 579
Chelonia mydas 702, 739
 Cheloniidae 698
Chelonoidis 770
Chelydra serpentina 153, 369–370
 Chilopoda 124–125
Chioninia 580
Chionodraco hamatus 234
Chirocephalus diaphanus 757–759
Chironius 770
 Chironomidae 634, 685
Chirostomias 394
Chonophorus 293, 297
Christinus 578
 chrysura 217
 sp. 217
Chrysemys 302, 586
 picta 770, 772
 Cichlidae 786
 Ciclantaceae 723
 Cladocera 759
Clariger 533
Clemmys guttata 179–190
Clidemia hirta 121
Cochliodon 494
Cochranella albomaculata 198
 Coleoptera 121, 124–127, 205, 207, 757–759
 Collembola 121–129, 205, 207
Coluber constrictor 446
 Colubridae 472–477, 580, 770
Compsaraia samueli 289
 Compsurini 243
Congoglanis 626, 629
 alula 626, 629
 howesi 626–630
 inga 626, 629
 sagitta 626, 629
Coniophanes andresensis 578
 Copepoda 685–686
 Copionodontinae 675
Cophixalus parkeri 729
Corallus 770
 hortulanus 472–477
Cordylus oelofseni 581
Coregonus 401, 405
Coris julis 233
 Corixidae 514
Corumbataia britskii 645–646
Corymbophanes 499, 524, 642
 andersoni 645
Cosmarium 514, 685
 Craugastoridae 171
Creagrutus mucipu 501
Cristatogobius lophius 533
Crocodylus
 johnstoni 702
 porosus 586
Crossodactylodes 683
Crotalus
 atrox 103
 cerastes 100–105, 119, 735
 horridus 103
 mittelli 103
 molossus 103
 oreganus 103
 pricei 119
Cryptoblepharus
 carnabyi 579
 egeriae 578
Cryptobranchus
 alleganiensis 150
 a. alleganiensis 150
 a. bishopi 150–159
Ctenocheirodon pristis 243–250
Ctenophorus clayi 581
Ctenosquamata 401, 405
Ctenotus 585
 alleni 581
 burbidgei 581
 ehmanni 581
 essington 578
 leonhardii 579
 Culicidae 685
 Curculionidae 205, 207
 Cyanophyceae 685–686
Cyclanthus 723
 sp. 723–724

- Cyclopoida 685, 757–759
Cyclodina alani 579
Cyclodomorphus
 casuarinae 578
 michaeli 578
Cynodontinae 618
Cynoscion regalis 142
Cyprinella
 monacha 167
 sp. 757
Cyprinodontidae 631
Cyprinodontiformes 762
Cyprinus carpio 169, 390
Cyrtocara sadleiri 579
Cyrtodactylus
 abrae 578
 eisenmanae 580
 eisenmani 580
 loriae 578
 sworderi 582
Cyrtopodion watsoni 582
Cyzicus grubei 757
- Dalatiidae* 766
Dallia 167
Danio rerio 776
Dasia 580
Dascyllus albisella 142
Dasyops 708
Dasythemis mincki 514
Delma 580
 haroldi 580
Delturus brevis 524
Dendrobates auratus 786
Dendrosaspis jamesoni 578
Dendropsophus cf. *microcephalus* 378
Dendrotriton 171
Dentex dentex 390
Derillissus 251–256
 altifrons 251, 253–254, 256
 krennobates 251, 253–256
 lombardii 251–253, 255–256
 nanus 251, 253–256
 vittager 251, 253, 255, 256
 vittiger 254, 256
Dermaptera 124–125
Dermatemys mawii 581
Dermochelys 772
 coriacea 739
Dermochylidae 698
Diacyclops 757
Diadophis 353
Diapoma speculiferum 611
Diaptomus cyaneus 757
Dibamus alfredi 581
Dicentrarchus labrax 777
Dicranopteris linearis 121
Dicroglossidae 108
Diffugia 685
Diplophidae 395, 405
Diplophos sp. 395, 401, 405
Diplopoda 124–125
Diplodactylus
 byrnei 581
 wombeyi 582
Diplura 205, 207
Diporiphora
 reginae 578
 winneckekei 582
Dipsas 197–202, 770
 albifrons 197
 indica 197
 nicholsi 197–202
 sp. 197–202
 variegata 197
Dipsosaurus dorsalis 102
Diptera 124–128, 205, 207, 757–759
Doumea 629
 alula 626
Doumeinae 626
Dreissena polymorpha 169
Dunhevedia crassa 757, 759
Duttaphrynus melanostictus 108
Dytiscidae 514
- Echiostoma* 395, 398, 405
 barbatum 396, 405
Egernia 130–134, 580
 coventryi 581
 cunninghami 130, 132
 douglasi 581
 frerei 130
 mcpheeii 579
 saxatilis 130, 132
 slateri 582
 stokesii 130, 132
 striolata 130–134
Elachistocleis surumu 378
Elassomatidae 167
Eleocharis palustris 757
Eleotris 533
 sandwichensis 297
Eleutherodactylidae 171
Eleutherodactylus 127–128
 coqui 121, 127–128, 729
 planirostris 121–129
Elimia potosiensis 371
Elseya jukesii 578
Emoia
 baudini 579
 kitcheneri 581
 reimschiisseli 581
Emydidae 301, 698–705
Emydoidea blandingii 188
Emydura
 australis 772
 macquarii 578, 772
Engystomops 507
Epactionotus 10
 bilineatus 645
Ephemeroptera 371–372, 634, 759
Epicrates 770
 cenchria 472–477
 c. cenchria 474–475
Epictia 43
 albipuncta 43, 46
 australis 46
 collaris 43

- columbi* 43
nasalis 43
peruviana 43
signata 43
striatula 43
tenella 43
undecimstriata 43
 Epictinae 37–48
 Epictini 37
Epidalea calamita 108, 111–112
 Epinephelidae 767–768
Epinephelus
 guttatus 134–135, 139, 142
 itajara 135, 142
 microdon 233
 morio 135, 142
 striatus 135, 141
Eremophilus 634
 mutisii 635
Eretmochelys imbricata 702, 739
Erimonax monachus 167
Eriophorum vaginatum 182, 184
Erythemis sp. 514
Erythrodiplax sp. 514
Euastrum 514
Eulamprus
 heatwolei 579
 martini 579
 Esocidae 167
Espadarana
 illex 197
 prosoblepon 197, 728
Etheostoma 432–439
 (Belophlox) 239
 chienense 437–438
 crossopterum 237, 432–439
 doration 239
 flabellare 239
 forbesi 237, 432–439
 fricksium 239
 inscriptum 237–242
 nigripinne 239, 432
 okaloosae 239
 rafinesquei 240–241
 rufilineatum 239
 squamiceps 239, 241, 432, 437–438
 tetrazonum 239
 (Ulocentra) 240
 Etmopteridae 766
Etmopterus
 pusillus 765
 robinsi 765
 virens 765
Eucalyptus 130
Euctenogobius 293
Eucypris virens 757
Eugerres mexicanus 163
Euglenophyta 685
Eunectes
 murinus 359
 notaues 359
Eupemphix 507
Eupogonesthes 394
Eurycheilichthys 10
 limulus 645
Eustomias 395, 398–399, 405
 bifilis 405
 filifer 405
 jimcraaddocki 160
 macrurus 405
 obscurus 396, 405
 sp. 398, 400, 406
Eutaeniichthys 533
Euteleostei 777
Eutenogobius 297
Exastilithoxus 547
 Fabaceae 205
Favia 217
Fejervarya
 cacrivora 108, 112
 limnocharis 108, 111–112
Festuca 204
Feylinia 580
Flagellostomias 395, 398–399, 405
 boureei 404
 Formicidae 121–129, 205, 207, 726
Fritziana goeldii 688
 Fundulidae 457
Fundulus 451
 albolineatus 167
 confluentus 456
 grandis 451, 456–457
 heteroclitus 451, 456–457
 jenkinsi 451–459
 luciae 451, 456
 parvipinnis 347, 456
 similis 451, 456–457
 Fungi 685–686
Galaxias 401
Galeichthys aguadulce 94
Galeorhinus galeus 765
Galeus
 antillensis 765
 cadenati 765
Gambusia affinis 786
Gasterosteus aculeatus 347
 Gastropoda 124–125
Geckonia chazaliae 578
Gehyra 580
Gekko 145, 147, 149
 badenii 578
 chinensis 145
 gecko 145, 149
 japonicus 145–149
 Geodipsadinae 580
Geomyersia 578
Geomys pinetus 706–713
Geophagus brasiliensis 523
Gerarda 580
 Ginglymostomatidae 766
Glandularia microphylla 205
Glaphyromorphus 578
 darwiniensis 578
 douglasi 581
Glauconia dimidiata 38, 40
Gloydus shedaoensis 119, 413, 416
Glyceria spicata 757
 Gobiesocidae 251–256

- Gobiesox*
lucayanus 256
punctulatus 256
Gobiidae 217, 527–534
Gobiiformes 359
Gobiodon histrio 217
Gongylomorphus bojerii 579
Gonocephalus sophiae 582
Gonostoma sp. 395, 401, 405
Gonostomatidae 395, 405
Gopherus 701
agassizii 222–228, 701
moraskai 222, 226
polyphemus 157, 701, 708
Guibemantis 648–662
aff. bicalcaratus 649, 656–657
aff. liber 650
albolineatus 648–649, 651, 657, 659
annulatus 648–649, 651, 657, 659–661
bicalcaratus 648–649, 651, 657, 659, 661
cf. bicalcaratus 661
cf. liber 650
cf. pulcher 649, 656–658
depressiceps 649, 657, 661
flavobrunneus 648–649, 651–652, 656–657, 659–661
(Guibemantis) 649–650, 656–657, 661
kathrinae 649, 657, 661
liber 648–662
(Pandanusicola) 648–662
pulcher 648–662
punctatus 648–649, 652, 657, 659
sp. 5 650
tasifotsy 648–662
timidus 649, 657, 661
tormieri 649, 657, 661
wattersoni 648–649, 652, 657, 659–661
Graciliscincus shonae 582
Grammatostomias 395, 398–399, 405
dentatus 405
Graptemys 301, 372, 699, 701
barbouri 301
caglei 301
ernsti 702
flavimaculata 305, 698–705
geographica 301, 367–374
n. delticola 702
oculifera 305, 702
ouachitensis 302, 304
o. ouachitensis 301–306
o. sabinensis 301–302, 304–305
pseudogeographica 301–305
p. ouachitensis 304
p. sabinensis 304
pulchra 301
versa 301
Gymnoamblyopus 527–534
Gymnogobius 533
Gymnophiona 168
Gymnotiformes 283–292, 401, 535–540
Gymnotocinclus 6, 11
anosteos 645
Gymnotus
albifrons 284
carapo 523
Hamptophryne 770
Harpacticoida 685
Harttia sp. 523
Hasemannia sp. 523
Hatcheria 634
macraei 635
Heleiporus australiacus 447
Heliconia 723
sp. 724–725, 727
Heliconiaceae 723
Hemiancistrus 547–553
medianus 552
Hemidactylus 147
frenatus 145, 148
mabouia 145, 148
turcicus 145
Hemidiaptomus maroccanus 757
Hemigrammus 611
Hemipsilichthys 524
gobio 524
nimius 524
papillatus 524
Hemiptera 124–125, 205, 207, 514, 757–759
Heptapteridae 675
Heterenchelyidae 484–493
Heterenchelys 491
biaggii 492
longus 491
macrurus 491
microphthalmus 491–492
Heterocheiron 243–244, 249
Heterodon platyrhinos 446
Heterodontidae 766
Heterophotus 394
ophistoma 405
Heteropriacanthus cruentatus 165
Heteroptera 124–125, 127
Hexapoda 205, 207, 686
Hippocampus 389–393
erectus 389–393
Hisonotus 266–275
armatus 272–273, 645
bockmanni 266–275
cf. depressicauda 273
cf. francirochai 273
cf. notatus 274
charrua 272
chromodontus 266–268, 271–273, 646
depressicauda 272–273
francirochai 273
hungry 272
insperatus 268, 272–273, 645–646
iota 272
laevior 274
leucofrenatus 274
leucophrys 272
luteofrenatus 266–268, 271–274, 646
megaloplax 272
montanus 272
nigricauda 272, 274
notatus 271–272
taimensis 272, 274
Hoffmannseggia minor 205

- Hollandichthys* 609, 611–613, 617–621
 multifasciatus 610, 612–614, 616–617, 619–622
Holocentridae 139
Homo sapiens 580
Homoptera 205, 207
Hoplias cf. *malabaricus* 523
Hoplobatrachus tigerinus 108, 112
Hoplocephalus
 bungaroides 119
 stephensi 415
Hoplodactylus
 rakiurae 578
 stephensi 578
Hyalinobatrachinae 729
Hyalinobatrachium 728–729
 aureoguttatum 722–731
 colymbiphyllum 730
 fleischmanni 728–730
 valerioi 727–730
Hydracarina 685–686
Hydrophis 277
 cantoris 277
 gracilis 277
 obscurus 277
 parviceps 276–282
Hyla
 altera 565
 avivoca 729
 chrysoscelis 729
 cinerea 729
 femoralis 729
 fuscomarginata 556, 565–566
 gratiosa 729
 meridionalis 757
 rubra
 r. altera 554, 556, 562, 565
 r. orientalis 554, 556
 squirella 729
Hylidae 108, 378, 770
Hylinae 171
Hylodes uai 523
Hymenoptera 121–129, 205, 207
Hypancistrus 547, 552
Hyphessobrycon 611
 bifasciatus 501
 heliacus 501
Hypomesus 401, 405
Hypoptopoma 6
 spectabile 646
Hypoptopomatinae 6, 10, 266–275, 498, 637, 641–643, 645
Hypoptopomatini 266
Hyporthodus 768
Hypostominae 498, 637–638, 642, 645
Hypostomini 642
Hypostomus 494–500, 638
 ancistroides 496–497, 499
 angipinnatus 496–497, 499
 aspilogaster 496–497, 499
 borellii 496–497, 499
 boulengeri 496–497, 499
 brevis 496–497, 499
 careopinnatus 494–500
 cochlodon 499
 commersoni 496–497, 499
 commersonii 645
 cordovae 497, 499
 denticulatus 497, 499
 derbyi 497, 499
 dlouhyi 497, 499
 fluviatilis 497, 499
 heraldoi 497, 499
 hermanni 497, 499
 isbrueckeri 497, 499
 laplatae 497, 499
 latifrons 497, 499
 levis 494–500
 margaritifer 499
 mutuae 497, 499
 nigromaculatus 497, 499
 peckoltoides 497, 499
 piratatu 497, 499
 regani 499
 roseopunctatus 499
 topavae 497, 499
 uruguayensis 497, 499
 variostictus 497, 499
Hypsiboas
 aff. *polytaenius* 514
 albomarginatus 564
 crepitans 378
 geographicus 378
 multifasciatus 378
 rosenbergi 729
 sp. 514
Hypsilyrus
 boydii 579
 godeffroyi 580
Ictaluridae 166
Ictalurus punctatus 88
Idiacanthus 395, 398, 405
 antrostomus 405
Iguania 12–22
Imantodes 770
Incilius nebulifer 106–114, 307–320
Insecta 124–125, 205, 207, 757–759
Isbrueckerichthys
 alipionis 274
 calvus 525
 epakmos 525
 saxicola 525
Isistius brasiliensis 765
Isogomphodon oxyrhynchus 765
Isopoda 121–129
Isoptera 205, 207
Isorineloricaria 494
Isthmohyla 171
Isurus
 oxyrinchus 788
 paucus 765
Juncus 451
Jupiaba pinnata 501
Kaloula pulchra 108, 112
Katrinus 578
Keratella 685

- Knodus*
 sp. 523
Kolpotocheiroidon 243
Kronichthys 6, 10–11
 subteres 274, 525, 646
- Labridae 217
Lacerta danfordi 581
Lacertaspis lepesmei 581
Lacertidae 580
Lachesis 770
Lagenophrys 685, 687
Lagenorhynchus acutus 160
Lagochila lacerum 167
Lamna nasus 788
Lamnidae 766
Lampetra minima 167
Lampropeltis
 getula getula 460–464
 triangulum 413, 415, 472–475
 t. micropholis 472–477
Lampropholis
 caligula 579
 guichenoti 580
Lapranthus sp. 121
Larrea tridentata 224
Lates niloticus 169
Lecane 685
Leiuperidae 378, 507–518
Leiurus berdmorei 579
Lentipes concolor 297
Lepadella 685
Lepidochelys
 kempii 701
 olivacea 738–747
Lepidodactylus mutahi 578
Lepidoptera 124–125, 205, 207
Lepidopus 34
Lepomis gibbosus 347
Leporacanthicus 547
Leporellus cf. *vittatus* 551
Leporinus
 desmotes 551
 tigrinus 551
Leptoancistrus 499, 524
Leptodactylidae 108, 378
Leptodactylus
 aff. *andreae* 378–379
 albilabris 108, 112
 bolivianus 378
 fuscus 378
 knudseni 378
 lineatus 378
 mystaceus 378
 podicipinus 378
 wagneri 378
Leptophis 770
Leptostomias 395, 398–399, 405
 gladiator 405
 sp. 405
Leptotyphlopidae 37–48
Leptotyphlops 37
 dimidiata 38
 dimidiatus 38, 40
 nursii 581
- Lerista*
 allanae 581
 bougainvillii 579
 christinae 579
 griffini 580
 haroldi 580
 humphriesi 581
 kendricki 581
 kennedyensis 578
 nevinae 580
 nicholli 580
Leuresthes tenuis 348
Leydigia sp. 757
Lialis 580
 burtoni 579
 jicari 581
Liasis 580
Libellulidae 514
Limnodynastes
 dumerilli 112
 peronei 112
Limulus polyphemus 422
Linderiella baetica 757–759
Linepithema humile 127
Liolaemidae 12–22
Liolaemus 12, 203, 786
 andinus 203
 alticolor 13
 bibronii 13
 capillitas 13, 18
 dorbignyi 203
 lutzae 203
 montanus 13, 17–18
 poecilochromus 203–210
 ruibali 17
 scrocchii 204
 tulkas 18
Liopholis
 kintorei 130
 whitii 130
Liparidae 257–265
Lipinia leptosoma 584
Lipnia 580
Lipoptericthys 524
Lithobates 168, 686
 areolatus 440–450
 capito 440
 catesbeianus 170, 424, 686
 clamitans 112, 194, 447
 (*Nenirana*) 440
 palustris 112, 440
 piapiens 109, 112, 321–330
 sevosus 440
 sp. 109
 sphenocephalus 312
 sylvaticus 109, 111–112, 312, 321–330, 424–431, 447
 virgatipes 109, 112
Lithogenes 163
Lithoxus 547
Litoria
 aurea 108, 111–112
 ewingii 109, 111–112
Lontra canadensis 151, 368, 373
Lophenchelys 484, 491

- fowleri* 484, 491
Loricaria 49–56
 apeltogaster 54–55
 cataphracta 54
 cf. *lentiginosa* 55
 cf. *simillima* 55
 coximensis 49–56
 holmbergi 49–56
 lentiginosa 54–55
 lundbergi 49–56
 nickeriensis 49–56
 parahybae 49–56
 piracicabae 55
 prolixa 54–55
 pumila 49–56
 simillima 54–55
 tucumanensis 54–55
Loricariidae 6–11, 49–56, 266–275, 494–500, 519–526, 547–553, 637–647, 675
Loricariinae 49
Loricarioidea 670, 675
Luciogobius 533
Lutjanidae 219
Lygisaurus sesbrauna 578
Lygodactylus 147
Lygosoma
 fernandi 578
 herberti 581
Mabuya 770
 stangeri 580
 stanjorgeri 580
Macadamia integrifolia 121
Macrochelys 772
Macropodus 347
Macropterus 552
Malacochersus tornieri 772
Malacosteus 394, 398, 405
 australis 396, 405
 niger 402, 405
Malacotraca 124–125
†*Maliamia gigas* 603–608
Mandarina 585
Mantella
 aurantiaca 649
 laevigata 648
Marantaceae 723
Maurolicus muelleri 405
Maxillopoda 757–759
†*Megacheiroidon* 243–244, 249
 unicus 244
Megadontognathus 289
Megalancistrus 547
Mehelya riggenbachi 581
Melanostomias 395, 398–399, 405
 biseriatus 405
 melanops 405
Melanosuchus niger 770
Menetia 580
 koshlandae 578
 sadlieri 578
Menidia 347
 menidia 341–350, 776–777
 peninsulae 346–348
Metacyclops sp. 757
Metrosideros polymorpha 121
Micralestes altus 483
Microhyla ornata 108, 112
Microhylidae 108, 378
Microlepidogaster 266, 645
 lophophanes 645
 perforatus 645
Micrurus 770
 averyi 581
Millipora 217
Mimagoniates rheocharis 611
Miscanthidium violaceum 383
Mixodiaptomus incrassatus 757
Moenkhausia pittieri 501
Monacanthus 579
Morelia 580
 macburnei 578
 mippughae 578
 oenpelliensis 578
Morethia 580
Moringua 489
Moringuidae 491
Mugilidae 167
Muraenidae 490
Musa 723, 725
 sp. 723–724
Musaceae 723
Mustela vision 151
Mustelus
 californicus 765
 canis 787
 felis 98–99
 higmani 765
Mycteroperca
 bonaci 135, 142
 interstitialis 137
 microlepis 141
 rosacea 142
 venenosa 135–144
 tigris 135–137, 141–142
Mylocheilus caurinus 347
Myoglanis 551
Myrocongridae 490

Nactus cheverti 578
Nannocharax
 cf. *fasciatus* 481
 parvus 481
Nannoplecostomus eleonora 637–647
Nanocheiroidon 243, 249
Nematobrycon 611
Nematocharax 611
 venustus 501
Nematogenys 675
Neolebias unifasciatus 481
Neonesthes 394–407
 capensis 406
Neoplecostominae 6, 499, 519, 646
Neoplecostomus 641
 microps 525, 646
 paranensis 274, 498
 sp. 523
Neoregelia cruenta 561, 566

- Neoteleostei 405
Neotoma
 floridana 708
 f. smalli 358
Nephrolepis sp. 121
Nerodia 358
 rhombifer 103
Nessia 580
 burtonii 579
 Neuroptera 124–125
Niveoscincus
 conventryi 581
 palfreymani 581
Nototriton 171
Novumbra 167
Nylanderia sp. 726

Odatia 580
 Odonata 514, 759
Odontamblyopus 527–534
Odontochelys 772
Odontostomias 394, 399
Odorrana 59, 65
 chapaensis 65, 69, 72
 geminata 65, 69, 72
 margaretae 59, 61–62
 schmackeri 59, 61–62
Oedipina 171
Oedogonium 514
Oligancistrus 552
 punctatissimus 551
Oligodon
 joynsoni 581
 perkinsi 581
Oligosarcus 611
 sp. 523
Oligosoma smithi 582
Ololygon 554
 altera 554, 556
Oncorhynchus 347
 aquabonita 167
 mykiss 151, 167, 395, 405
Oncorhynchus rhodurus 234
Ondatra zibethicus 358
 Ophichthidae 279, 489
Ophidia 580
Ophisichthys 491
 dubius 491
 Opiliones 726
Opistomias 395, 398–399, 405
 mitsuii 406
Orconectes
 longidigitus 151
 neglectus 151
 punctimanus 151
Oreochromis
 mossambicus 386
 niloticus 234, 777
Orestias 631
Orthemis sp. 514
 Orthoptera 124–125, 205, 207
Orthosternarchus 289
Oryzias 776
 latipes 776

Osmerus 401, 405
 sp. 151
Ostariophys 637
Osteichthyes 762
Osteocephalus 770
Osteoglossum bicirrhosum 770
Ostracoda 757–759
 †*Ostrea meunieri* 604–605
Otocinclus 266, 641
 bororo 646
 caxarari 646
 depressicauda 273
 flexilis 645
 francirochai 273
 hasemani 646
 mangaba 646
 tapirape 645–646
Otothyriini 266, 271
Otothyris
 lophophanes 274, 645–646
 rostrata 646
Otothyropsis marapoama 274
 Oxynotidae 766
Oxyrhopus 770
 erdicii 581
Oxyurichthys 533
 ophthalmonema 533
 visayanus 533

 †*Pachyamia* 607
Pachydactylus turneri 582
Pachyseris 217
Pachystomias 395, 398, 405
 microdon 396, 406
Pagrus major 234
Pailsus pailsei 578
Panaque 547
Pandanus 648–662
Pantodactylus nicefori 579
Panturichthys 484, 488, 490–491
 fowleri 484–493
 isognathus 489–491
 longus 490–491
 mauritanicus 489–492
Papuascincus stanleyanus 578
 Paragoniatinae 611
Parancistrus nudiventris 551
Paraneetroplus maculicauda 786
Parapistocalamus hedigeri 579
Parapteronotus hasemani 289
Pareiorhaphis 519–526, 641
 cameroni 525
 eurycephalus 519–526
 nasuta 519–526
 nudulus 519–526
 parmula 525
 proskynita 519–524
 scutula 519–526
 vestigipinnis 499, 519–526
Pareiorhina 498, 524, 641
 carrancas 499, 524–525
 rudolphi 274, 525, 646
Parotocinclus 10, 266, 271, 638
 amazonensis 645–646

- aripuanensis* 646
bahiensis 646
britskii 646
collinsae 645–646
eppleyi 646
jumbo 645
longirostris 646
maculicauda 645
muriaensis 646
polyochrus 646
prata 6, 10–11, 274, 645
Pauropoda 124–125
Peckoltia 547–553
 bachi 547–553
 compta 552
 feldbergae 547–553
 oligospila 547–553
 sabaji 547–553
 snethlageae 552
 vittata 552
Pelobates cultripes 757
Pelophylax
 klepton esculentus 194
 perezi 109
Pennatula 253
Pennisetum clandestinum 121
Percidae 432–439, 714
Perciformes 714
Percina 719
 nigrofasciata 714–721
Peromyscus 708
Petromyzon marinus 169
Phacus 685
Phalloceros sp. 523
Phasmahyla jandaia 523
Pheidole megacephala 127
Phelsuma 147, 580, 584–585
 madagascariensis kochi 581
Phenacobrycon 621
Phenacogrammus aurantiacus 481
Philodendron 723, 725
 sp. 724
Philodryas carbonelli 579
Phormidium 514
Phosichthyidae 395, 405
Photonectes 395, 398, 405
 braueri 406
 sp. 400, 406
Photostomias 394, 398–399, 405
 guernei 403, 406
 sp. 395, 397–398, 406
Phrynocephalus grumgrzimailoi 581
Phyllodytes 683
 gyrinaethes 688
Phyllomedusa
 hypochondrialis 378, 729
 sauvagii 729
Phyllomedusinae 171
Phyllurus championae 579
Phymaturus 12–22, 203
 agilis 13
 alica huense 13–15
 antofagastensis 13, 15, 18
 calcogaster 13–14, 18
 castillensis 13–14, 18
 ceii 13–14, 18
 cf. *palluma* 18
 cf. *punae* 18
 darwini 13–15
 desuetus 13–14
 dorsimaculatus 13, 15, 18
 etheridgei 13–14, 18
 excelsus 13–14, 18
 extrilidus 13, 16–18
 felixi 13–14, 19
 gynechlomus 13
 indistinctus 13–14, 19
 laurenti 13, 15, 19
 mallimaccii 13, 15, 19
 manuelae 13–14, 19
 maulense 13–15
 nevadoi 13–14, 19
 paihuanense 13–16, 19
 palluma 12–22
 patagonicus 12–14, 19
 payunia 13–14, 19
 punae 13, 15–16, 18–19
 querque 13, 15, 19
 roigorum 13, 15, 19
 somuncurensis 13–14, 19
 sp. 20
 spectabilis 13–14, 20
 spurcus 13–14, 20
 tenebrosus 13–14, 20
 verdugo 13, 15, 20
 videlai 13–14, 20
 vociferator 13–15
 zapalensis 13–14, 20
Physalaemus
 aguirrei 514–515
 albifrons 507–508, 514–516
 albonatus 514–516
 angrensis 515
 atlanticus 516
 barrioi 507–518
 biligonigerus 514–515
 bokermanni 516
 camacam 516
 centralis 514, 516
 combrei 516
 crombiei 514
 cuqui 514
 cuvieri 507–508, 514–516
 deimaticus 507–508, 514
 enesefae 678–682
 ephippifer 378, 514
 erikae 514–515
 evangelistai 507, 514
 feioi 515
 fernandezae 514–516
 gracilis 507–518
 henseli 508, 514–515
 henselii 507, 514
 jordanensis 507, 514, 516
 laterstriga 515
 lisei 507, 514, 516
 maculiventris 516
 marmoratus 514–515

- maximus* 515
moreirae 514, 516
olfersi 515
olfersii 507–508, 514
orophilus 515
palluma 12–22
pustulosus 507, 678–682
riograndensis 514–515
santafecinus 514, 516
santafeicus 515
signifer 507, 514–515
soaresi 515
Picea glauca 332
Pimelodus
 albidus 86
 spixii 81, 86
Pimephales notatus 438
Pinus
 clausa 706
 contorta 332
 echinata 706
 elliottii 706
 palustris 706–713
 taeda 706
Pituophis
 catenifer 710
 melanoleucus 706, 710
 m. lodgini 710
 m. mugitus 706–713
Placobdella cryptobranchii 154
Plasmodium 766
Platemys 586
Plecoptera 372
Plecostomus 494
 scaphyiceps 499
 variostictus 499
Plectropomus leopardus 219
Plesiodypsas 197
Plesioptopoma 6–11
 curvidens 6–11
Plethodon 753
 cinereus 748–755
Plethodontidae 171
Pleurodeles waltl 756–761
Pleuronectiformes 541–546
Pocillopora 217–218
Podocnemis 770
 vogli 578
Podocopida 757–759
Poduromorpha 685
Poecilia latipinna 386
Pogona
 henrylawsoni 579
 vitticeps 416
Pogonopoma obscurum 645
Polypedes megacephalus 108, 112
Polypogon maritimus maritimus 757
Pomacentridae 217
Pomacentrus
 amboinensis 217
 moluccensis 217
 partitus 142
Populus tremuloides 182, 332
Porites 217–218
Porotergus ellisi 284
Potamopyrgus antipodarum 169
Potamorrhaphis 663–669
 eigenmanni 663, 665–668
 guianensis 663, 665–669
 labiatus 663–668
 petersi 663, 665–669
Prasinohaema 584
Precathorops 77
Premnas biaculeatus 217
Pristimantis 770
Proablepharus reginae 578
Procambarus sp. 422
Proctoporus unsaaca 578
Procyon lotor 151, 184, 446
Protacanthopterygii 401
Protozoa 685–686
Pseudacris
 crucifer 109, 112
 kalmi 109, 112
 maculata 321–330
 regilla 687, 754
 sp. 109
 triseriata 112
Pseudancistrus 552
 cf. *barbatus* 551
Pseudecheneis weigeli 578
Pseudechis
 porphyriacus 413
 weigeli 580
Pseudemoia baudini 579
Pseudemys concinna 370
Pseudocarcharhidae 766
Pseudochalceus 611, 619
Pseudocorynopoma 621
Pseudocrenilabrus
 multicolor 382–388
 m. victoriae 383
Pseudonaja
 elliotti 578
 gowi 578
 ingrami 580
Pseudorinelepis 499
 genibarbis 499
Pseudoscorpiones 124–125
Pseudothecadactylus lindneri 580
Pseudotocinclus 6, 10–11
 juquiai 11
 parahybae 11
 tietensis 274, 645
Pseudotothyris obtusa 274
Pseudotrypauchen 527–534
Pseudotylosurus 663
Psidium cattleianum 121
Psocoptera 121–129
Ptenopus garrulus garrulus 145, 147–148
Pterobrycon 621
Pterygoplichthys 552
Ptyodactylus hasselquisti 148
Pygidium
 areolatum 635
 schmidtii 635
Pygopus schraderi 580
Python

- bivittatus* 170, 358
regius 359
reticulatus 359
schneideri 579
Pythonichthys 486–487, 489–491
asodes 486, 489, 491
macrurus 486, 491
microphthalmus 491–492
sanguineus 490–492

Quercus 706
falcata 707
laurifolia 707
nigra 707
virginiana 707

Rachoviscus 609, 611–613, 617–621
crassiceps 610, 612–617, 620–622
graciliceps 609, 612–614, 616–618, 620–622
Ramphotyphlops
becki 578
exocoeti 578
robertsi 580
sylvia 578
tovelli 582
Rana 168, 690
catesbeiana 695
lessonae 429
muscosa 154
pipiens 695
pretiosa 157
temporaria 110, 112, 427, 447
Ranidae 109–110
Rattus 358
Rena 37–48
affinis 40, 43, 46
bressoni 37
dimidiata 37–48
dissecta 37
myopica 37
unguirostris 40, 43, 46
Retama monosperma 757
Retroculus xinguensis 551
Rhabdosargus haffara 234
Rhacophoridae 108, 110
Rhadinesthes 394–407
decimus 406
Rhaebo guttatus 378
Rhamdia quelen 163
Rhantus
(Rhantus) 514
sp. 514
Rhinelepini 642
Rhinelepis 498
aspera 499
Rhinella 770
cf. merianae 378
icterica 514
marina 108, 111–112, 378
Rhinichthys
atratus 786
cobitis 167
obtus 786
Rhinophis fergusonianus 581
Rhizoprionodon porosus 765

Rhizosomichthys 631
totae 635
Rhoadsiinae 611
Rhyacichthys aspro 294–296, 299
Rhypticus
nigromaculatus 30, 32
subbifrenatus 30
Riopa 580
Ristella 580
rurkii 580
Rocio octofasciata 164
Rotifera 685–686
Rubus idaeus strigosus 182, 184
Rypticus 23–36
bistrispinus 34
bornoi 24–25, 34
carpenteri 24–30, 32–35
maculatus 34
randalli 34
saponaceus 24–25, 34
sp. 24–25, 28
subbifrenatus 23–36

Sachatamia illex 728
Saiphos 580
Salmininae 618
Salmo trutta 151
Salmonidae 405
Saltuarius
kateae 581
swaini 579
Sapotaceae 723–725
Saprosincus 578
challengeri 578
galli 579
rosei 579
Sauria 580
Scaphiophis raffreyi 581
Scaphiopus holbrookii 447
Scaridae 229–236
Scarthyla 770
Scarus ferrugineus 229–236
Sceloporus 586
occidentalis 702
pilsbryi 579
scalaris 786
thayerii 579
Scenodesmus 685
Schizolecis guntheri 274
Schroederichthys maculatus 765
Sciaenidae 167
Scinax 683–689, 770
acuminatus 556, 559, 563, 565–566
aff. alter 562
alter 554–569
altera 556, 561, 564
alterna 554
alterus 556
auratus 556, 562
blairi 556, 559, 563, 565
cabralensis 556
camposseabrai 556
castroviejoi 556
catharinae 554

- cf. alter* 565
chiquitanus 556
cretatus 562
crospodospilus 556
cruentomimus 556, 559–560, 563–565
cuspidatus 556, 559–560, 562–566
duartei 556
elaeochrous 556
eurydice 556, 559–560, 563–566
funereus 556, 559–560, 563–565
fuscovarius 556, 559, 563–565, 567
granulatus 556, 559–560, 563–565, 567
hayii 556
ictericus 556, 559–560, 563–565
imbegue 554–569
iquitorum 556, 559–560, 563–565
karenanneae 556, 559, 563, 565
lindsayi 556, 559, 563, 565
littoreus 683–688
machadoi 523
manriquei 556
maracaya 556
oreites 556
pachycrus 556, 560, 563, 565, 567
perereca 556
perpusillus 683–689
quinquefasciatus 556
rostratus 554, 556
ruber 378, 554–569
similis 556, 560, 563–565, 567
sp. 565–566
sp. aff. alter 565
sp. I 562, 564
sp. II 565–566
squalirostris 514
staufferi 554, 556
tigrinus 556
tymbamirim 554–569
uruguayus 554, 556
x-signatus 556
Scincus philbyi 579
Scirpus maritimus 757
Scobinancistrus 552
Scolopacidae 670, 675
Scoloplax 670–677
baileyi 670–675
baskini 670, 672–677
dicra 672–676
distolothrix 672–676
dolicholophia 672–676
empousa 672–676
Scopelarchus 402, 405
Scopelosaurus craddocki 160
Scorpaeniformes 257–265
Scyliorhinidae 766
Scymnodon 766
Serpentes 37–48
Serranidae 23–36, 135–144, 217–218, 768
Serrapinnus 243–244, 249
heterodon 244, 523
Serrasalminae 619
Siagonodon 43
acutirostris 43
borrichianus 43, 46
cupinensis 43
septemstriatus 43
Sibon 197–202
annulatus 197–202
argus 197–202
longifrenis 197–202
nebulatus 197–202
Sibynomorphus 197
Sicyopterus
lagocephalus 297
stimpsoni 297
Sigara *sp.* 514, 757
Sigmodon 358
Siluriformes 6–11, 49–56, 77–97, 266–275, 494–500, 519–526, 547–553, 626, 637, 670, 675–677
Silvinichthys 631
bortayro 631, 634–635
leoncitensis 631, 634
mendozensis 631, 634–635
sp. 631
sp. A 635
sp. B 635
Simoselaps roperi 578
Siphophis worontzowi 582
Sistrurus
catenatus 119, 735
c. catenatus 156
miliaris 358
Solea
fonsecensis 543
panamensis 541–546
Solifugae 205, 207
Somniosidae 766
Somniosus 766
Spartina 451
Spectracanthicus 547
Sphaenorhynchus 770
Sphaerodactylus 145
clenchi 581
rosaurae 578
Sphagneticola trilobata 121
Sphagnum 182
Sphenodon
punctatus 586
Sphenomorphus
coxi 581
kuehnei 580
Sphyrna 34, 788
Sphyrnidae 766
Spinicaudata 757–759
Spinomantis
brunae 654
nussbaumi 654
Spintherobolus 243–244, 249
Squalidae 766
Squaliforma 494
Squaloliparis dentatus 264
Squalus 766
suckleyi 765
Squatina
californica 765
mexicana 765
Squatinae 766
Stenodus leucichthys 166

- Stenogobius hawaiiensis* 297
Stenostoma 44
 dimidiata 37
 dimidiatum 38
Stenotaphrum secundatum 121
Sternarchogiton 289
Sternarchorhynchus 289
Sternarchorhamphus 289
Sternarchorhynchus
 higuchii 290
 jaimeii 290
Sternarchus
 brasiliensis 284
 leptorhynchus 284
 macrostomus 284
 mariae 284
 rostratus 284
 spurrellii 284
Sternoptychidae 395, 405
Sternoptyx 395
 sp. 395, 405
Sternopygidae 551
Sternorrhyncha 124–126
Sternotherus odoratus 370, 701, 770, 772
Stevardiinae 619–620, 622
Stomias 394–407
 boa 396, 398, 402–403, 405–406
 colubrinus 404, 406
 lampropeltis 399, 403, 406
Stomiidae 394–407
Stomiiformes 405
Stonogobius 533
Storeria dekayi 578
Strabomantidae 171
Streptocephalus torvicornis 757–759
Strophurus
 jeanae 580
 krisalys 581
 wilsoni 580
Stylophora 217
Synodontidae 219

Tachisurus
 agassizii 94
 nuchalis 78
 spixii 83
Tachysurus
 agassizii 91
 nuchalis 78
 spixii 83
Tactostoma 395, 398–399, 405
 macropus 406
Taenioides
 jacksoni 527–534
 limicola 527–534
Tantilla 770
Tanymastix sp. 757–758
Teleocichla
 centrarchus 551
 monogramma 551
Teleostei 23–36, 631–636, 211–221, 229–236, 251–256,
 394–407, 786
Tenebrionidae 205, 207
Teratohyla pulverata 728

Terrarana 171
Testudinidae 698
Testudo
 klienmanni 156
Tetragnatha 585
Tetragonopterinae 611, 619
Tetragonopterus carvalhoi 501
Thalassoma
 hardwicke 217
 jansenii 217
 lunare 217
Thammophis 356–357, 587
 elegans 103
 e. vagrans 334
 sirtalis 446
 s. sirtalis 446
Thysanactis 395, 398–399, 405
 dentex 406
Thysanoptera 124–126, 128, 205, 207
Tiaroga cobitis 167
Tiliqua 580
Tomicodon
 cryptus 256
 fasciatus 256
 reitzei 256
 rupestris 256
Tomiyamichthy 533
Trachelomonas 514
Trachemys
 scripta 701–702
 s. elegans 367–374
Trachops cirrhosus 729
Trachycephalus 770
 typhonius 378
Trachylepis
 brauni 579
 hemmingi 581
Triakidae 98–99
Triakis
 felis 98
 semifasciata 98–99, 766
 semifasciatum 98
Tricheilostoma 37–48
 anthracinum 44, 46
 brasiliensis 44, 46
 brevissimum 44, 46
 dimidiatum 38, 43–44, 46
 dugandi 43–44, 46
 fuliginosum 44, 46
 guayaquilensis 43–44, 46–47
 jani 39–45, 47
 joshuai 43–44, 47
 koppesi 44, 47
 macrolepis 44, 47
 nicefori 43–44, 47
 salgueiroi 44, 47
Trichogenes 675
Trichomycteridae 675
Trichomycterinae 635
Trichomycterus 631–636
 aguarague 632, 635
 alterus 631–632, 634, 635
 areolatum 635
 areolatus 631, 635

- barbouri* 631, 635
belensis 631–632, 635
borellii 631, 635
boylei 631–635
catamarcensis 635
corduensis 631–636
duellmani 634–635
gracilis 635
heterodontus 631
hualco 631, 635
megantoni 635
minus 631–636
nigricans 634
pseudosilvinichthys 631, 635
punctulatus 635
ramosus 631–633, 635
riojanus 631
rivulatus 635
roigi 634–635
schmidtii 635
sp. 523
spgazzinii 631, 635
therma 634
tiraquae 634
yuska 631, 635
Trichoptera 371–372
Trigonomolampa 394
Trinectes 541–546
 fimbriatus 541, 543
 fluviatilis 541, 543
 fonsecensis 541–546
 hubbsbollinger 541–546
 inscriptus 541, 543
 maculatus 541, 543
 microphthalmus 541, 543
 opercularis 541, 543
 paulistanus 541, 543
 xanthurus 541, 543
Triturus vulgaris 336
Tropidechis sadlieri 578
Tropidodipsas 197
Tropidonophis
 mairii 580
 parkeri 580
Tropidophorus baviensis 579
Trypanosoma 766
Trypauchenopsis 527–534
 intermedia 527–534
Turbellaria 757
Typhlopidae 579
Typhlops kraali 581
Typhlosaurus
 cregoi 581
 veris 578
Ubidia magdalenensis 284
Umbra 167
Umbridae 167
Urodela 168
Uta stansburiana 593–602
Vaccinium corymbosum 182
Varanus rosenbergi 580
Verbenaceae 205
Vermicella snelli 582
†*Vidalamia* 607
 catalunica 604
Vidalamiinae 604
Vidalamiini 603–608
Vinciguerra
 poweriae 405
 sp. 395, 405
Vipera berus 472
Viperidae 100–105
Vriesea friburgensis 561
Waglerophis merremi 472
Watawata 494
Xantusia riversiana 578
Xenentodon 663
Xenochrophis 586
Xenopus 690–691, 695
 laevis 690–697
Xiphophorus 776
 multilineatus 786
Zeus 34
Zygnemaphyceae 685

SUBJECT INDEX

Copeia 2012, Nos. 1–4

- ACTIVITY**, *Crotalus cerastes* (seasonal aphasia) 100–105; *Bitis schneideri* (sexual seasonal movement) 115–120; *Eleutherodactylus planirostris* (feeding, diet) 121–129; *Mycteroperca venenosa* (spawning aggregations; sound production; communication) 135–144; *Gekko japonicus* (acoustic/social behavior) 145–149; *Clemmys guttata* (habitat selection; movement) 179–190; *Acris crepitans* (dispersal; movement) 191–196; *Chaetodon (auriga, aureofasciatus, rainfordi, plebius)*, *Chelmon rostratus* (larval settlement) 211–221; *Gopherus agassizii* (nesting) 222–228; *Ambystoma macrodactylum* (movement patterns) 331–340; *Bothrops insularis* (thermoregulation) 408–418; *Etheostoma (forbesi, crossoplerum)* (nesting) 432–439; *Lampropeltis g. getula* (diel patterns) 460–464; *Basiliscus plumifrons* (budget variation) 465–471; Boidae, Colubridae (intraspecific recognition) 472–477; *Physalaemus barroii* (calling parameters; intrageneric comparison) 507–518; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.)* (advertisement calls) 554–569; *Pituophis melanoleucus mugitus* (movement; home range) 706–713; *Bitis schneideri* (movement; seasonality) 732–737.
- AESTIVATION**, *Clemmys guttata* (habitat conditions) 179–190.
- AGE**, *Liolaemus poecilochromus* (diet, juveniles/adults) 203–210.
- AGGREGATIONS**, *Egernia striolata* (tree-dwelling populations; kinship) 130–134; *Mycteroperca venenosa* (spawning; sound production) 135–144.
- AVOIDANCE**, *Lithobates sylvaticus* (olfactory cues; migration) 424–431.
- AWARDS**, W. F. Smith-Vaniz, R. D. Semlitsch, H. R. Mushinsky, M. B. Thompson, P. H. Skelton 1–5.
- BEHAVIOR**, *Phymaturus (extrilidus n. sp.)* (parental care) 12–22; *Crotalus cerastes* (seasonal aphasia) 100–105; *Egernia striolata* (aggregations) 130–134; *Mycteroperca venenosa* (aggregations; sound production; spawning) 135–144; *Gekko japonicus* (acoustic; social) 145–149; *Clemmys guttata* (flooding effects: habitat selection; home range; movement) 179–190; *Acris crepitans* (dispersal; movement) 191–196; *Chaetodon (auriga, aureofasciatus, rainfordi, plebius)*, *Chelmon rostratus* (larval settlement) 211–221; *Gopherus agassizii* (nesting; anthropogenic variables) 222–228; *Graptemys geographica* (basking site) 367–374; *Pseudocrenilabrus multicolor* (mouth-brooding) 382–388; *Bothrops insularis* (thermoregulation; seasonal variation) 408–418; *Lithobates sylvaticus* (juvenile orientation) 424–431; *Etheostoma (forbesi, crossoplerum)* (nesting) 432–439; *Lithobates areolatus* (migration; burrow philopatry) 440–450; *Fundulus jenkinsi* (reproduction) 451–459; *Basiliscus plumifrons* (activity budget: seasonal variation, ontogenetic) 465–471; Boidae, Colubridae (tongue flicking) 472–477; *Panturichthys fowleri* (feeding, habitat) 484–493; *Peckoltia (feldbergae n. sp.)* (rheophilic; nocturnal foraging) 547–553; *Guibemantis (tasifotsy n. sp.)* (breeding site) 648–662; *Physalaemus pustulosus* (calls; acoustic environment) 678–682; *Scinax (littoreus, perpusillus)* (tadpole feeding) 683–689; *Hyalinobatrachium aureoguttatum* (male clutch attendance) 722–731; *Lepidochelys olivacea* (hatchling emergence) 738–747; *Plethodon cinereus* (color change) 748–755; *Pleurodeles wahl* (newt diet) 756–761.
- BIODIVERSITY**, *Phymaturus* (33 spp.; increase) 12–22; *Amolops (marmoratus sp. complex)* (Myanmar cryptic spp.) 57–76; *Cathorops* (review W. Atl. spp.) 77–97; Characidae (tribe Cheirodontini: *Clenocheiroidon n. gen.*) 243–250; *Apterionotus* (nominal spp., table) 283–292; *Awaous* (Pacific; evolutionary lineages) 293–300; anurans (riparian/non-riparian communities) 375–381; *Scinax (alter sp. complex)* (new species identified) 554–569; *Uta stansburiana* (reproductive isolation; speciation) 593–602; *Congoglanis (howesi n. sp.)* (Congo R. basin endemic) 626–630; *Potamorhaphis (labiatus n. sp.)* 663–669; *Scoloplax (baileyi n. sp., baskini)* (Rio Negro basin increase) 670–677.
- BIOGEOGRAPHY**, *Plesiopoma (curvidens n. gen., n. sp.)* (type loc. only: trib. upper Rio Sao Francisco, Braz.) 6–11; *Phymaturus* (clade range; vicariance) 12–22; *Tricheilostoma (dimidiatum n. comb.)* (SE Venez. record doubtful), (*jani n. sp.*) (limited, Espinhaço Range, Braz.) 37–48; *Loricaria (coximensis n. sp.)* (Braz., near power plant) 49–56; *Eleutherodactylus planirostris* (Hawaiian introduction) 121–129; *Egernia striolata* (juv. delay of dispersal) 130–134; *Apterionotus* (nominal spp.: distribution, habitat) 283–292; *Awaous (guamensis, stamineus)* (Hawaiian Is.) 293–300; *Bathyaethiops* (range extension) 478–483; *Astyanacinus (yariguies n. sp.)* (1st record, Rio Magdalena basin) 501–506; *Trypauchenopsis intermedia* (wide-spread; collection difficulties) 527–534.
- BURROWS**, *Lithobates areolatus* (fidelity; movement) 440–450.
- CALLS**, *Gekko japonicus* (acoustic behavior; intra-/intersexual communication; spectrograms) 145–149; *Physalaemus barroii* (advertisement: audiospectrum, oscillogram; 26 spp., call parameters) 507–518; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.)* (advertisement; audiospectrum) 554–569; *Guibemantis (tasifotsy n. sp.)* (sonogram, oscillogram) 648–662; *Physalaemus pustulosus* (acoustic environment; evoked vocal response) 678–682; *Hyalinobatrachium aureoguttatum* (microhabitat selection) 722–731.
- CANNIBALISM**, *Pseudocrenilabrus multicolor* (filial; mouth brooders) 382–388.
- CHEMORECEPTION**, *Lithobates sylvaticus* (orientation cues) 424–431; Boidae, Colubridae (intraspecific recognition) 472–477.
- COLONIZATION**, *Inciilius nebulifer* (invasive, disturbed habitats) 307–320.
- COLORATION**, *Plesiopoma (curvidens n. gen., n. sp.)* (in alcohol; b&w photo.) 6–11; *Phymaturus (extrilidus n. sp.)* (in life/alcohol; b&w photos.; color fig.; variation) 12–22; *Rypticus (carpenteri n. sp., subbifrenatus)* (in life; color photos.; comparison; diagnostic) 23–36; *Tricheilostoma (dimidiatum n. comb.)* (in alcohol), (*jani n. sp.*) (in life/alcohol; b&w photos.) 37–48; *Loricaria (coximensis n. sp.)* (in alcohol; b&w photo.) 49–56; *Amolops (marmoratus, afghanus, indoburmanensis n. sp.)* (in preserv.; b&w photos.) 57–76; *Cathorops (nuchalis, spixii, wayui n. sp.)* (in life; b&w photos.) 77–97; *Mycteroperca venenosa* (color phases; reprod. chgs.; color photos.) 135–144; *Clenocheiroidon (pristis n. gen., n. sp.)* (in life; color photo.) 243–250; *Derilissus (lombardii n. sp.)* (in life/alcohol; color photos.), (*altifrons, kremnobates, vittiger, nanus*) 251–256; *Careproctus (hyperciscus n. sp., kamikawai n. sp., ovigerus)* (in life/preserv.; color photo.) 257–265; *Hisonotus (bockmanni n. sp.)* (unique pattern wrt congeners) 266–275; *Hydrophis parviceps* (counter-coloration; b&w/color photos.) 276–282; *Apterionotus (lindalvae n. sp.)* (in alcohol; diagnostic) 283–292; *Bathyaethiops (atercrinis n. sp.)* (in alcohol; b&w/color photos.) 478–483; *Panturichthys fowleri* (in life; b&w photos.) 484–493; *Hypostomus (careopinnatus n. sp.)* (in alcohol; b&w photos.) 494–500; *Astyanacinus (yariguies n. sp.)* (in alcohol; unique pattern; b&w photo.) 501–506; *Physalaemus barroii* (in life/preserv.; b&w photo./fig.) 507–518; *Pareiorhaphis (proskynita n. sp.)* (in life/alcohol; unique pattern; color photo.) 519–526; *Trypauchenopsis intermedia* (in life/alcohol; b&w figs.) 527–534; *Adontosternarchus (duartei n. sp.)* (in alcohol; b&w photo.) 535–540; *Trinectes (hubbsbolinger n. sp.)* (in preserv.; color photos.) 541–546; *Peckoltia (feldbergae n. sp.)* (in life/alcohol; unique pattern; color photos.) 547–553; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.)* (in life; b&w photos./figs.) 554–569; *Uta stansburiana* (speciation; habitat component) 593–602; *Congoglanis (howesi n. sp.)* (in alcohol; b&w photo.) 626–630; *Trichomycterus (minus n. sp.)* (in alcohol; diagnostic; photo.) 631–636; *Nannoplecostomus (eleonorae n. gen., n. sp.)* (in alcohol; photo.) 637–647; *Guibemantis (tasifotsy n. sp.)* (in life/alcohol; photos.) 648–662; *Potamorhaphis (labiatus n. sp.)* (in life/alcohol; photos.) 663–669; *Scoloplax (baileyi n. sp.)* (in life/alcohol; photos.) 670–677; *Plethodon cinereus* (pigmentation change in captivity) 748–755.

- COMMUNICATION.** *Mycteroperca venenosa* (social/reproductive behavior; oscillo-/spectrograms) 135–144; *Gekko japonicus* (intra-/intersexual) 145–149; Boidae, Colubridae (chemoreception & intraspecific recognition) 472–477; *Physalaemus barrooi* (advertisement call) 507–518; *Guibemantis* (*tasifotsy* n. sp.) (calls) 648–662; *Physalaemus pustulosus* (evoked vocal response) 678–682.
- COMMUNITY.** *Egernia striolata* (kinship; tree-dwell populations) 130–134; *Graptemys geographica*, *Trachemys scripta elegans* (structure; population shifts; habitat alteration) 367–374; anurans (comparison, riparian/non-riparian) 375–381; *Bathyaethiops* (*atercrinis* n. sp.) (structure) 478–483; *Pareiorhaphis* (*proskynita* n. sp.) (structure) 519–526; *Peckoltia* (*feldbergae* n. sp.) (structure) 547–553.
- COMPETITION.** *Chaetodon* (*auriga*, *aureofasciatus*, *rainfordi*, *plebicus*), *Chelmon rostratus* (larval settlement; non-predatory residents) 211–221; *Hollandichthys*, *Rachoviscus* (sperm competitiveness) 609–625.
- CONSERVATION.** *Bitis schneideri* (habitat fragmentation) 115–120; *Mycteroperca venenosa* (over-fishing) 135–144; *Cryptobranchus alleganiensis bishopi* (captive-reared survival in wild; habitat) 150–159; *Sibon*, *Dipsas* (dietary implications) 197–202; *Gopherus agassizii* (nest-site, anthropogenic variables) 222–228; *Hydrophis parviceps* (endangered status) 276–282; *Awaous stamineus* (poss. Haw. endemic) 293–300; *Lithobates* (*pipiens*, *sylvaticus*), *Pseudacris maculata*, *Anaxyrus* (*cognatus*, *woodhousii*) (ecological niche theory; habitat suitability) 321–330; *Ambystoma macrodactylum* (road mortality reduction) 331–340; *Graptemys geographica* (habitat degradation; basking site loss) 367–374; anurans (riparian habitat & species diversity) 375–381; *Fundulus jenkinsi* (spawning habitat) 451–459; *Physalaemus barrooi* (State & IUCN status) 507–518; *Scoloplax* (*baileyi* n. sp.) (widespread; protected areas) 670–677; *Lepidochelys olivacea* (hatchling emergence; sand temperature) 738–747.
- DEEP-SEA.** *Careproctus* (*lycopersicus* n. sp., *kamikawai* n. sp., *ovigerus*) (No. Pacific; bathy/benthic, 2300–3000 m) 257–265; Stomiidae (osteology; dorsal gill arches; ontogeny) 394–407.
- DENTITION.** *Plesiopoma* (*curvidens* n. gen., n. sp.) (premax., photo.) 6–11; *Cathorops* (*nuchalis*, *spixii*, *wayuu* n. sp.) (male/female tooth plates, fig.) 77–97; *Ctenochirodon* (*pristis* n. gen., n. sp.) (premax., max., dentary; photo.) 243–250; *Careproctus* (*lycopersicus* n. sp., *kamikawai* n. sp., *ovigerus*) (jaw teeth) 257–265; *Hydrophis parviceps* (counts) 276–282; *Bathyaethiops* (*atercrinis* n. sp.) (premax., dentary teeth, fig.) 478–483; *Panturichthys fowleri* (upper/lower jaws, photos; variation) 484–493; *Asyanacinus* (*yariguies* n. sp.) (jaw teeth; fig.) 501–506; *Trichomycterus* (*minus* n. sp.) (diagnostic) 631–636; *Scoloplax* (*baileyi* n. sp.) (new unique tooth morphology) 670–677.
- DEVELOPMENT.** *Incilius nebulifer* (salinity effects) 106–114; *Atherinops affinis*, *Menidia menidia* (latitudinal variation, Jordan's Rule; sex determination) 341–350; *Xenopus laevis* (tadpole, environmental factors, respiration) 690–697.
- DIET.** *Phymaturus* (*extrilidus* n. sp.) (herbivory) 12–22; *Loricaria* (*coximensis* n. sp.) (seeds) 49–56; *Eleutherodactylus planirostris* (native/non-native invert. spp.) 121–129; *Sibon* (*longifrenis*, *argus*, *annulatus*, *nebulatus*), *Dipsas* (*nicholsi*, sp.) (gut analysis) 197–202; *Liolaemus poecilochromus* (herbivory) 203–210; *Pareiorhaphis* (*proskynita* n. sp.) (mainly algae) 519–526; *Peckoltia* (*feldbergae* n. sp.) (herbivore) 547–553; *Scinax* (*tymbamirim* n. sp.) (arthropods) 554–569; *Scinax* (*littoreus*, *perpusillus*) (selectivity; seasonality; categories) 683–689; *Pleurodeles waltl* (new feeding; zooplankton) 756–761.
- DIGESTION.** *Alligator mississippiensis* (rate; hard-body prey) 419–423.
- DISEASE.** *Cryptobranchus alleganiensis bishopi* (released captive-reared, susceptibility) 150–159.
- DISPERSAL.** *Bitis schneideri* (home range; seasonal movement) 115–120; *Egernia striolata* (delay by juveniles) 130–134; *Acris crepitans* (movement, habitat-type effect) 191–196; *Awaous* (amphidromus gobioid) 293–300; *Graptemys* (*o. ouachitensis*) (Pleistocene glacial refugia) 301–306; *Incilius nebulifer* (invasive; disturbed habitats) 307–320; *Lithobates sylvaticus* (orientation cues) 424–431; *Lithobates areolatus* (juvenile migration) 440–450.
- DISTRIBUTION.** *Plesiopoma* (*curvidens* n. gen., n. sp.) (type loc. only; trib. Rio Sao Francisco, Braz.; map) 6–11; *Phymaturus* (*extrilidus* n. sp.) (type loc. only; San Juan Prov., Arg.; map), (table, 33 recognized spp.) 12–22; *Rypticus* (*carpenteri* n. sp.: Bermuda & Caribbean), (*subbifrenatus*: Caribbean, parts W. Afr. coast), (map) 23–36; *Tricheilostoma* (*jani* n. sp.: limited Espinhaço Range, Braz.), (*dimidiatum* n. comb.: Braz. Amaz. lowlands, Guiana, Surinam Centro-Roraima), (map) 37–48; *Loricaria* (*coximensis* n. sp.) (Rio Paraguay basin, Braz.; map) 49–56; *Amolops* (*marmoratus*: Mon & Shan states & no. Tanintharyi Div.), (*afghanus*: no. Kachin state), (*panhai*: record for Myanmar), (*indoburmanensis* n. sp.) (Chin & Rakhine states & Sagaing Div.), (map) 57–76; *Cathorops* (*nuchalis*: Venez. to Guyana), (*spixii*: Amaz. R. to Rio de Janeiro), (*wayuu* n. sp.: no. Colombia to Venez.), (map; incl. C. [agassizii, arenatus, mapale], *Arius* [*variolosus*, *nigricans*]) 77–97; *Bitis schneideri* (endemic, so. Afr. Succulent Karoo) 115–120; *Eleutherodactylus planirostris* (Haw. Is. introduction) 121–129; *Mycteroperca venenosa* (Caribbean; map) 135–144; *Sibon* (so. Cent. Amer. to no. So. Amer.), *Dipsas* (Mex. to cent. So. Amer.) 197–202; *Ctenochirodon* (*pristis* n. gen., n. sp.) (Rio Tocantins, Braz.; map) 243–250; *Derilissus* (*lombardii* n. sp.) (holotype only: Exumas) 251–256; *Careproctus* (*lycopersicus* n. sp., *kamikawai* n. sp., *ovigerus*) (Bering Sea, NE Pacific; deep-sea; map) 257–265; *Hisonotus* (*bockmanni* n. sp.) (type loc. only: Rio Tapajós, Amaz. Basin; map) 266–275; *Hydrophis parviceps* (so. Vietnam; prob. endemic) 276–282; *Apternotus* (nominal spp.; table), (*lindae* n. sp.: Rio Uatumã, Amaz.; map) 283–292; *Awaous* (*guamensis*, *stamineus*) (Guam, Haw. Is.; map) 293–300; *Graptemys* (*o. ouachitensis*: thru-out Mississippi R. drainage), (*o. sabinensis*: Sabine R. only), (*pseudogeographica*: middle Mississippi R. drainage) 301–306; *Anaxyrus fowleri*, *Incilius nebulifer* (range decrease/increase) 307–320; *Etheostoma forbesi* (endemic; Caneys Fork R. syst., TN) 432–439; *Bathyaethiops* (range extension), (*atercrinis* n. sp.) (type loc. only: Louessé R.; map) 478–483; *Panturichthys fowleri* (east. Mediterranean; map) 484–493; *Hypostomus* (*careopinatus* n. sp.) (type loc. only: Rio Taquari drainage, Braz.; map) 494–500; *Asyanacinus* (*yariguies* n. sp.) (tribs., Rio Cascajales, Colombia; map) 501–506; *Physalaemus barrooi* (endemic, Atl. Forest, SE Braz.) 507–518; *Pareiorhaphis* (*proskynita* n. sp.) (upper Ribeirão Caraca, Braz.; map) 519–526; *Trypauchenopsis intermedia* (Indo-W. Pac.) 527–534; *Adontosternarchus* (*duartei* n. sp.) (Rio Purus basin, Braz.; map) 535–540; *Trinectes* (*hubbbsollinger* n. sp.) (limited, tribs., Pac. slope, Colombia) 541–546; *Peckoltia* (*feldbergae* n. sp.) (Rio Xingu basin only; map) 547–553; *Scinax* (*alter*, *imbegue* n. sp., *tymbamirim* n. sp.) (coastal, east. Braz.; map) 554–569; *Congoglanis* (*howesi* n. sp.) (type loc. only: Kasai R. endemic; map) 626–630; *Trichomycterus* (*minus* n. sp.) (endemic, Andean stream, Argentina) 631–636; *Nannoplecostomus* (*eleonoreae* n. gen., n. sp.) (tribs., upper Rio Tocantins basin, Braz.; map) 637–647; *Guibemantis* (*tasifotsy* n. sp.) (regional endemic, SE Madagascar; map) 648–662; *Potamorhaphis* (*labiatus* n. sp.) (west. Amaz. basin; map) 663–669; *Scoloplax* (*baileyi* n. sp.) (tribs., Rio Unini; map), *S. baskini* (range extension) 670–677.
- ECOLOGY.** *Plesiopoma* (*curvidens* n. gen., n. sp.) (distribution; habitat; osteology; sexual dimorphism) 6–11; *Phymaturus* (*extrilidus* n. sp.) (distribution; habitat; diet; color variation; basking/body temps.; parental care) 12–22; *Rypticus* (*carpenteri* n. sp., *subbifrenatus*) (distribution; habitat) 23–36; *Tricheilostoma* (*jani* n. sp., *dimidiatum* n. comb.) (distribution; habitat; sexual dimorphism) 37–48; *Loricaria* (*coximensis* n. sp.) (distribution; habitat; diet; sexual dimorphism; paedomorphism) 49–56; *Amolops* (*marmoratus*, *afghanus*, *panhai*, *indoburmanensis* n. sp.) (distribution; sexual dimorphism) 57–76; *Cathorops* (*nuchalis*, *spixii*, *wayuu* n. sp.) (distribution; sexual dimorphism; coloration; habitat) 77–97; *Crotalus cerastes* (reproduction; feeding) 100–105; *Incilius nebulifer* (larval survival; salinity effects) 106–114; *Bitis schneideri* (distribution; habitat; male/female seasonal movement; conservation) 115–120; *Eleutherodactylus planirostris* (diet) 121–129; *Egernia striolata* (aggregations; juvenile dispersal; kinship) 130–134; *Mycteroperca venenosa* (aggregations; social/reproduction behavior; sound production; color phases) 135–144; *Gekko japonicus* (behavior; communication; sound) 145–149; *Cryptobranchus alleganiensis bishopi* (survival/body condition wrt habitat) 150–159; *Clemmys guttata* (habitat selection; home range; movement patterns; sex ratio; aestivation) 179–190; *Acris crepitans* (dispersal; movement; habitat

- type) 191–196; *Sibon* (*longifrenis argus*, *annulatus*, *nebulatus*), *Dipsas* (*nicholsi*, sp.) (diet; distribution; natural history) 197–202; *Liolaemus poecilochromus* (diet; ontogeny; sexual dimorphism; natural history; parasites) 203–210; *Chaetodon* (*auriga*, *aureofasciatus*, *rainfordi*, *plebius*), *Chelmon rostratus* (larval settlement; habitat selection; movement) 211–221; *Gopherus agassizii* (nesting; hatching success; reproduction) 222–228; *Etheostoma inscriptum* (reproduction; sex ratio; sexual dimorphism; GSI; fecundity; life history) 237–242; *Clenocheirodon* (*pristis* n. gen., n. sp.) (distribution; sexual dimorphism; habitat) 243–250; *Derilissus* (*lombardii* n. sp.) (habitat) 251–256; *Careproctus* (*lycopersicus* n. sp., *amikawai* n. sp., *ovigerus*) (distribution; habitat; sexual dimorphism) 266–275; *Hydrophis parviceps* (distribution; endemism; feeding; sexual dimorphism) 276–282; *Apteronomus* (*lindalvae* n. sp.) (distribution; habitat; endemism; feeding; sexual dimorphism) 283–292; *Awaous* (*guamensis*, *stamineus*, *ocellaris*) (distribution; life history; evolutionary lineages) 293–300; *Anaxyrus fowleri* (range decrease), *Incilius nebulifer* (range increase) 307–320; *Lithobates* (*pipiens*, *sylvaticus*), *Pseudacris maculata*, *Anaxyrus* (*cognatus*, *woodhousii*) (habitat) 321–330; *Ambystoma macrodactylum* (breeding site migration) 331–340; *Atherinops affinis*, *Menidia menidia* (Jordan's Rule, latitudinal variation: vertebral number, sexual determination) 341–350; *Graptemys* (*geographica*, *Trachemys scripta elegans*) (behavior; community interactions; habitat alteration; population size; sex ratio; feeding) 367–374; anurans (habitat & species diversity) 375–381; *Pseudocrenilabrus multicolor* (mouth brooding; gill size/position & respiration; filial cannibalism) 382–388; *Hippocampus erectus* (sexual dimorphism) 389–393; *Bothrops insularis* (behavior; thermoregulation; seasonal variation) 408–418; *Alligator mississippiensis* (feeding; digestion rate) 419–423; *Lithobates sylvaticus* (juvenile orientation; olfactory cues) 424–431; *Etheostoma* (*forbesi*, *crossotermum*) (nesting; hybridization; reproduction) 432–439; *Lithobates areolatus* (migration; movement/forays; burrow philopatry; mortality/survivorship) 440–450; *Fundulus jenkinsi* (reproduction; spawning season; oogenesis) 451–459; *Lampropeltis g. getula* (diel activity patterns) 460–464; *Basiliscus plumifrons* (activity budget; habitat use) 465–471; Boidae (4 spp.), Colubridae (1 sp.) (chemoreception; intraspecific recognition) 472–477; *Bathyaethiops* (*atercrinis* n. sp.) (habitat; distribution) 478–483; *Panturichthys fowleri* (habitat; distribution; inferred feeding/habit) 484–493; *Hypostomus* (*careopinnatus* n. sp.) (distribution; habitat) 494–500; *Aspynacianus* (*yariguies* n. sp.) (distribution) 501–506; *Physalaemus barrioi* (distribution; habitat; advertisement call; tadpole morphology; karyotype) 507–518; *Pareiorhaphis* (*proskynita* n. sp.) (distribution; habitat; diet; community; sexual dimorphism) 519–526; *Trypauchenopsis intermedia* (distribution; habitat; sexual dimorphism) 527–534; *Adontosternarchus* (*duartei* n. sp.) (distribution; habitat) 535–540; *Trinectes* (*hubbbsollinger* n. sp.) (distribution) 541–546; *Peckoltia* (*jeldbergae* n. sp.) (distribution; habitat; diet; sexual dimorphism; community) 547–553; *Scinax* (*alter*, *imbegue* n. sp., *tymbamirim* n. sp.) (distribution; habitat; calls; diet; breeding season; fecundity) 554–569; *Uta stansburiana* (reproductive isolation; habitat; color morphs; speciation) 593–602; †*Vidalmiini* (newfossil specimen; marine environment; Eocene; figs.) 603–608; *Congoglanis* (*howesi* n. sp.) (distribution; habitat) 626–630; *Trichomycterus* (*minus* n. sp.) (distribution; habitat) 631–636; *Nannoplecostomus* (*eleonora* n. gen., n. sp.) (distribution; habitat; sexual dimorphism) 637–647; *Guibemantis* (*tasifotsy* n. sp.) (distribution; habitat; breeding site; calls) 648–662; *Potamorhaphis* (*labiatus* n. sp.) (distribution; habitat; sexual dimorphism) 663–669; *Scoloplax* (*baileyi* n. sp.) (distribution; habitat), *baskini* (range extension) 670–677; *Physalaemus pustulosus* (calls; acoustic environment) 678–682; *Scinax* (*litoreus*, *perpusillus*) (tadpole diet; trophic niche) 683–689; *Xenopus laevis* (tadpole growth & development; environmental factors) 690–697; *Graptemys flavimaculata* (stress response; corticosterone levels) 698–705; *Pituophis melanoleucus mugitus* (home range; habitat selection; refugia) 706–713; *Percina nigrofasciata* (reproductive life history; seasonal/temporal variation) 714–721; *Hyalinobatrachium aureoguttatum* (male clutch attendance; microhabitat; calling) 722–731; *Bitis schneideri* (movement) 732–737; *Lepidochelys olivacea* (hatchling emergence temperature; sex ratio) 738–747; *Plethodon cinereus* (color change in captivity) 748–755; *Pleurodeles waltli* (newt feeding; prey choice) 756–761.
- EGGS, *Gopherus agassizii* (clutch size; hatching success) 222–228; *Careproctus* (*lycopersicus* n. sp., *amikawai* n. sp., *ovigerus*) (sizes) 257–265; *Etheostoma* (*forbesi*, *crossotermum*) (clutch size; chronology) 432–439; *Scinax* (*tymbamirim* n. sp.) (clutch size) 554–569; *Uta stansburiana* (clutch size; reproductive isolation) 593–602; *Percina nigrofasciata* (size; clutch size) 714–721; *Hyalinobatrachium aureoguttatum* (clutch size) 722–731.
- ENDANGERED SPECIES, *Loricaria* (*coximensis* n. sp.) (human activity) 49–56; *Mycteroperca venenosa* (over-fishing) 135–144; *Cryptobranchus alleganiensis bishopi* (release/survival captive-reared) 150–159; *Hydrophis parviceps* (IUCN Red List) 276–282; *Apteronomus* (*lindalvae* n. sp.) (dam construction) 283–292; *Etheostoma forbesi* (restricted habitat; hybridization) 432–439; *Fundulus jenkinsi* (state/federal status) 451–459; *Physalaemus barrioi* (State & IUCN status) 507–518; *Peckoltia* (*jeldbergae* n. sp.) (aquarium trade; dam construction) 547–553; *Trichomycterus* (*minus* n. sp.) (mining run-off) 631–636.
- ENDEMIC/ENDEMISM, *Bitis schneideri* (Succulent Karoo, so. Afr.; habitat fragmentation/conservation mgt.) 115–120; *Hydrophis parviceps* (prob. status, so. Vietnam) 276–282; *Awaous stamineus* (poss., Haw. Is.) 293–300; *Etheostoma forbesi* (restricted, Caneey Fork R. syst., TN) 432–439; *Physalaemus barrioi* (Atl. Forest, SE Brazil; life history) 507–518; *Congoglanis* (*howesi* n. sp.) (Kasai R.) 626–630; *Trichomycterus* (*minus* n. sp.) (type loc. only: Altiplano stream, Arg.) 631–636; *Guibemantis* (*tasifotsy* n. sp.) (regional, SE Madagascar) 648–662.
- EVOLUTION, *Egernia striolata* (social behavior; aggregations) 130–134; *Awaous* (*guamensis*, *stamineus*) (evolutionary lineages) 293–300; *Uta stansburiana* (reproductive isolation; speciation process) 593–602; †*Vidalmiini* (marine amiids; persistence, Cretaceous–Paleocene boundary) 603–608.
- FECUNDITY, *Etheostoma inscriptum* (multiple spawnings) 237–242; *Etheostoma* (*forbesi*, *crossotermum*) (clutch size) 432–439; *Scinax* (*tymbamirim* n. sp.) (clutch size) 554–569; *Uta stansburiana* (clutch size; reproductive isolation; population crosses) 593–602; *Percina nigrofasciata* (clutch size; seasonal/temporal variation) 714–721; *Hyalinobatrachium aureoguttatum* (clutch size; male clutch attendance) 722–731.
- FEEDING, *Crotalus cerastes* (seasonal aphasia) 100–105; *Eleutherodactylus planirostris* (diet; native/non-native inverts.) 121–129; *Sibon* (*longifrenis*, *argus*, *annulatus*, *nebulatus*), *Dipsas* (*nicholsi*, sp.) (diet; goo-eaters) 197–202; *Liolaemus poecilochromus* (diet; ontogenetic shift, variation) 203–210; *Hydrophis parviceps* (prob. snake eels) 276–282; *Graptemys* (*geographica*) (dietary specialist; habitat alteration) 367–374; *Alligator mississippiensis* (digestion rate) 419–423; *Panturichthys fowleri* (inferred diet) 484–493; *Scinax* (*litoreus*, *perpusillus*) (tadpole oral morphology) 683–689; *Pleurodeles waltli* (newt prey choice) 756–761.
- FORAGING, *Basiliscus plumifrons* (activity budget, sexual/ontogenetic) 465–471.
- FOSSIL, †*Vidalmiini* (Eocene; marine) 603–608.
- GENETICS, *Rypticus* (*carpenteri* n. sp., *subbifrenatus*) (mtDNA) 23–36; *Amolops* (*marmoratus*, *afghanus*, *panhai*, *indoburmanensis* n. sp.) (mtDNA; cryptic spp.; consensus tree) 57–76; *Egernia striolata* (μsat. DNA; aggregation kinship) 130–134; *Hydrophis parviceps* (mtDNA) 276–282; *Awaous guamensis* (mtDNA; evolutionary lineages; phylogenetic analysis) 293–300; *Graptemys* (*o. ouachitensis*, *o. sabinensis*, *pseudogeographica*) (mtDNA; variation; haplotypes; ML tree) 301–306; *Menidia menidia* (GSD, genetic sex determination) 341–350; *Ambystoma* (cross-species microsatellite primers), *A. annulatum* (polymorphic loci) 570–577; *Uta stansburiana* (speciation; reproductive isolating mechanism; population crosses) 593–602; *Guibemantis* (*tasifotsy* n. sp., *pulcher*, *liber*) (mtDNA; tree, fig.) 648–662.
- GEOGRAPHIC LOCALITIES, Angola, *Congoglanis* (*howesi* n. sp.) 626–630. Antigua, *Rypticus subbifrenatus* 23–36. Argentina, *Phymaturus* (*extridius* n. sp., *palluma* gr., *patagonicus* gr.) 12–22; *Liolaemus poecilochromus* 203–210; *Trichomycterus* (*minus* n. sp.) 631–636.

- Arkansas, *Graptemys o. ouachitensis* 301–306.
 Atlantic Ocean, (NW) *Menidia menidia* 341–350.
 Australia, (NSW) *Egernia striolata* 130–134; (GBR) *Chaetodon (auriga, aureofasciatus, rainfordi, plebius)*, *Chelmon rostratus* 211–221.
 Bahamas, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36; *Derilissus (lombardii n. sp.)* 251–256.
 Belize, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Bermuda, *Rypticus (carpenteri n. sp.)* 23–36.
 Brazil, *Plesiopoma (curvidens n. gen., n. sp.)* 6–11; *Rypticus subbifrenatus* 23–36; *Tricheilostoma (jani n. sp.)* 37–48; *Loricaria (coximensis n. sp.)* 49–56; *Cathorops spixii* 77–97; *Ctenocheirodon (pristis n. gen., n. sp.)* 243–250; *Hisonotus (bockmanni n. sp.)* 266–275; *Apteronotus (lindalvae n. sp.)* 283–292; riparian anurans 375–381; *Bothrops insularis* 408–418; *Hypostomus (careopinatus n. sp.)* 494–500; *Physalaemus barrovi* 507–518; *Pareiorhaphis (proskynita n. sp.)* 519–526; (Amaz.) *Adontosternarchus (duartei n. sp.)* 535–540; *Peckoltia (feldbergae n. sp.)* 547–553; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.)* 554–569; *Hollandichthys (Rachoviscus 609–625; Nannoplecostomus (leonora n. gen., n. sp.)* 637–647; *Potamorhaphis (labiatus n. sp.)* 663–669; *Scoloplax (baileyi n. sp., baskini)* 670–677; *Scinax (littoreus, perpusillus)* 683–689.
 California, *Gopherus agassizii* 222–228; *Uta stansburiana* 593–602.
 Caroline Islands, *Trypauchenopsis intermedia* 527–534.
 Canada, (Ont.) *Clemmys guttata* 179–190; (Alb.) *Ambystoma macrodactylum* 331–340.
 Chile, *Phymaturus (palluma gr.)* 12–22.
 Colombia, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36; *Cathorops (wayuu n. sp.)* 77–97; *Astyanacinus (yariguies n. sp.)* 501–506; *Trinectes (hubbsollinger n. sp.)* 541–546; *Hyalinobatrachium aureoguttatum* 722–731.
 Congo, Republic of, *Bathyaethiops (atercrinis n. sp.)* 478–483.
 Costa Rica, *Basiliscus plumifrons* 465–471.
 Cuba, *Rypticus subbifrenatus* 23–36.
 Curaçao, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Dominica, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Dominican Republic, *Rypticus subbifrenatus* 23–36.
 Ecuador, Boa (*c. constrictor, c. imperator*), *Corallus hortolanus*, *Epicratis encrinura*, *Lampropeltis triangulum micropholis* 472–477.
 Equatorial Guinea, *Rypticus subbifrenatus* 23–36.
 Fernando Poo, *Rypticus subbifrenatus* 23–36.
 Florida, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36; *Hippocampus erectus* 389–393; *Percina nigrofasciata* 714–721.
 Georgia, *Lampropeltis g. getula* 460–464; *Pituophis melanoleucus mugitus* 706–713.
 Grand Cayman, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Grenadines, *Rypticus subbifrenatus* 23–36.
 Guam, *Awaous (guamensis, ocellaris)* 293–300.
 Guiana, *Tricheilostoma (dimidiatum n. comb.)* 37–48.
 Guyana, *Cathorops (nuchalis, wayuu n. sp.)* 77–97.
 Haiti, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Hawaii, *Eleutherodactylus planirostris* 121–129; *Awaous (guamensis, stamineus)* 293–300.
 Honduras, *Rypticus subbifrenatus* 23–36.
 Illinois, *Graptemys (o. ouachitensis, pseudogeographica)* 301–306.
 Indiana, *Lithobates areolatus* 440–450.
 Indonesia, *Trypauchenopsis intermedia* 527–534; *Lepidochelys olivacea* 738–747.
 Jamaica, *Rypticus subbifrenatus* 23–36.
 Japan, *Gekko japonicus* 145–149.
 Kansas, *Graptemys (o. ouachitensis, pseudogeographica)* 301–306.
 Kentucky, *Graptemys o. ouachitensis* 301–306.
 Louisiana, *Anaxyrus fowleri*, *Iniclus nebulifer* 307–320; *Graptemys (o. ouachitensis, o. sabinensis)* 301–306.
 Madagascar, *Guibemantis (tasifotsy n. sp., pulcher, liber)* 648–662.
 Maine, *Lithobates sylvaticus* 424–431.
 Marianas Islands, *Trypauchenopsis intermedia* 527–534.
 Massachusetts, *Plethodon cinereus* 748–755.
 Mediterranean Sea, (E) *Panturichthys fowleri* 484–493.
 Mexico, (Carib.) *Rypticus subbifrenatus* 23–36.
 Minnesota, *Graptemys (o. ouachitensis, pseudogeographica)* 301–306.
 Mississippi, *Fundulus jenkinsi* 451–459; *Graptemys flavimaculata* 698–705.
 Missouri, *Cryptobranchus alleganiensis bishopi* 150–159; *Graptemys (o. ouachitensis, pseudogeographica)* 301–306; *Graptemys geographica*, *Trachemys scripta elegans* 367–374.
 Myanmar, *Amolops (marmoratus, afghanus, panhai, indoburmanensis n. sp.)* 57–76.
 Navassa, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Nigeria, *Rypticus subbifrenatus* 23–36.
 North Dakota, *Lithobates (pipiens sylvaticus)*, *Pseudacris maculata*, *Anaxyrus (cognatus, woodhousii)* 321–330.
 Ohio, *Acris crepitans* 191–196; *Graptemys o. ouachitensis* 301–306; *Plethodon cinereus* 748–755.
 Pacific Ocean, (NE) *Careproctus (lycopersicus n. sp., kamikawai n. sp., ovigerus)* 257–265; (NE) *Atherinops affinis* 341–350.
 Panama, *Rypticus subbifrenatus* 23–36; *Sibon (longifrenis, argus, annulatus nebulatus)*, *Dipsas (nicholsi, sp.)* 197–202.
 Pennsylvania, *Plethodon cinereus* 748–755.
 Peru, *Potamorhaphis (labiatus n. sp.)* 663–669.
 Philippines, *Trypauchenopsis intermedia* 527–534.
 Puerto Rico, *Rypticus subbifrenatus* 23–36; *Mycteroperca venenosa* 135–144.
 Ryukyu Islands, *Trypauchenopsis intermedia* 527–534.
 Saba, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Saudi Arabia, (Red Sea) *Scarus ferrugineus* 229–236.
 Senegal, *Vidalmiini* 603–608.
 South Africa, *Bitis schneideri* 115–120, 732–737; *Trypauchenopsis intermedia* 527–534.
 South Carolina, *Etheostoma inscriptum* 237–242.
 Spain, *Pleurodeles waltl* 756–761.
 Surinam, *Tricheilostoma (dimidiatum n. comb.)* 37–48.
 Tennessee, *Etheostoma (forbesi, crossopterus)* 432–439.
 Tobago, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Turks & Caicos, *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 Uganda, *Pseudocrenilabrus multicolor* 382–388.
 Venezuela, *Rypticus subbifrenatus* 23–36; *Cathorops (nuchalis, wayuu n. sp.)* 77–97.
 Vietnam, *Hydrophis parviceps* 276–282.
 Virgin Islands (U.S.), *Rypticus (carpenteri n. sp., subbifrenatus)* 23–36.
 West Virginia, *Graptemys o. ouachitensis* 301–306.
 Wisconsin, *Graptemys o. ouachitensis* 301–306.
 GONOSOMATIC INDEX (GSI), *Etheostoma inscriptum* (peak) 237–242; *Fundulus jenkinsi* (spawning season) 451–459.
 GROWTH, *Hippocampus erectus* (allometry) 389–393; *Xenopus laevis* (tadpole, environmental factors, respiration) 690–697.
 HABITAT, *Plesiopoma (curvidens n. gen., n. sp.)* (type loc.; sm. stream) 6–11; *Phymaturus (extrilidus n. sp.)* (fissured rock outcrops) 12–22; *Rypticus (carpenteri n. sp., subbifrenatus)* (similar coral/rock, different depths) 23–36; *Tricheilostoma (jani n. sp.)* (variable in transition areas; photos.), *(dimidiatum n. comb.)* (dry regions in transition areas) 37–48; *Loricaria (coximensis n. sp.)* (riffles) 49–56; *Cathorops (nuchalis, spixii, wayuu n. sp.)* (coastal, into brackish water) 77–97; *Bitis schneideri* (coastal sand dunes) 115–120; *Mycteroperca venenosa* (aggregation site, shelfbreak, 20–50 m depth) 135–144; *Cryptobranchus alleganiensis bishopi* (suitable release sites for captive-reared) 150–159; *Clemmys guttata* (flood effects; selection; movement patterns; home range) 179–190; *Acris crepitans* (effect on movement/dispersal) 191–196; *Liolaemus poelochromus* (cold-climate Andes & diet) 203–210; *Chaetodon (auriga, aureofasciatus, rainfordi, plebius)*, *Chelmon rostratus* (selection/larval settlement) 211–221; *Gopherus agassizii* (anthropogenic changes & nesting) 222–228; *Ctenocheirodon (pristis n. gen., n. sp.)* (holotype: clear water) 243–250; *Derilissus (lombardii n. sp.)* (deep mesophotic reef; photo.) 251–256; *Careproctus (lycopersicus n. sp., kamikawai n. sp., ovigerus)* (deep-sea, 2300–3000 m; prob. rock ledge) 257–265; *Hisonotus (bockmanni n. sp.)* (type loc. only: sandbank) 266–275; *Hydrophis parviceps* (marine; unknown) 276–282; *Apteronotus (lindalvae n. sp.)* (rheophilic), (nominal spp. list) 283–292; *Anaxyrus fowleri*, *Iniclus nebulifer* (disturbed urban areas; invasive sp.) 307–320; *Lithobates (pipiens, sylvaticus)*, *Pseudacris maculata*, *Anaxyrus (cognatus, woodhousii)* (grasslands, wetlands; ecological niche theory) 321–330; *Ambystoma macrodactylum* (breeding site migrations & soil moisture) 331–340; *Graptemys geographica*,

- Trachemys scripta elegans* (degradation, population size) 367–374; anurans (riparian, Amazon forest) 375–381; *Pseudocrenilabrus multicolor* (hypoxic & mouth brooders) 382–388; *Bothrops insularis* (usage wrt thermoregulation) 408–418; *Etheostoma (forbesi, crossopterum)* (nesting sites) 432–439; *Fundulus jenkinsi* (salt marsh; spawning sites) 451–459; *Lampropeltis g. getula* (longleaf pine forest) 460–464; *Basiliscus plumifrons* (behavior/seasonal usage; activity budget) 465–471; *Bathyaethiops (atercrinis n. sp.)* (type loc. only; photo.) 478–483; *Panturichthys fowleri* (muddy bottoms, 30–120 m depth) 484–493; *Hypostomus (careopinnatus n. sp.)* (type loc.: sm. stream) 494–500; *Physalaemus barrioi* (highlands, Atl. Forest; endemic) 507–518; *Pareiorhaphis (proskynita n. sp.)* (alt. >1000 m; physical characters) 519–526; *Trypauchenopsis intermedia* (mud burrows, estuarine) 527–534; *Adontosternarchus (duartei n. sp.)* (white-water river) 535–540; *Peckoltia (feldbergae n. sp.)* (shallow rapids) 547–553; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.)* (arboreal/shrub) 554–569; *Uta stansburiana* (reproductive isolating mechanism btwn. populations) 593–602; †*Vidalmiini* (Eocene; marine deposit; map) 603–608; *Congoglanis (howesi n. sp.)* (rapids) 626–630; *Trichomycterus (minus n. sp.)* (thermal water; 2318 m elev.) 631–636; *Nannoplecostomus (eleonora n. gen., n. sp.)* (sm. tributaries, karst region) 637–647; *Guibemantis (tasifotsy n. sp.)* (vegetation, lowland swamps), *G. pulcher* (*Pandanus* leaf axils) 648–662; *Potamorrhaphis (labiatus n. sp.)* (sm. streams, lagoons) 663–669; *Scoloplax (bailei n. sp.)* (blackwater streams) 670–677; *Scinax (littoreus, perpusillus)* (bromeliad pools) 683–689; *Pituophis melanoleucus mugitus* (longleaf pine; selection) 706–713; *Hyalinobatrachium aureoguttatum* (microhabitat selection) 722–731; *Bitis schneideri* (Succulent Karoo) 732–737; *Pleurodeles waltl* (vernal pools) 756–761.
- HATCHLING**, *Lepidochelys olivacea* (emergence sand temperature & success; sex ratio) 738–747.
- HISTOLOGY**, *Scarus ferrugineus* (ovaries; oogenesis) 229–236; *Hollandichthys*, *Rachoviscus* (sperm) 609–625.
- HOME RANGE**, *Bitis schneideri* (male/female, seasonal movement) 115–120; *Clemmys guttata* (flood. effects) 179–190; *Lithobates areolatus* (burrow philopatry; travel distance) 440–450; *Pituophis melanoleucus mugitus* (size; selection; variation) 706–713.
- HYBRIDIZATION/HYBRIDS**, *Etheostoma (forbesi, crossopterum)* (nesting activity; chronology) 432–439.
- JORDAN'S RULE**, *Atherinops affinis*, *Menidia menidia* (latitudinal variation & vertebral number) 341–350.
- KINSHIP**, *Egernia striolata* (tree-dwelling populations) 130–134.
- LARVAE**, *Incilius nebulifer* (salinity levels & effects on hatching/metamorphosis) 106–114; *Chaetodon (auriga, aureofasciatus rainfordi, plebius)*, *Chelmon rostratus* (settlement; habitat selection; movement) 211–221; Stomiidae (ontogeny; dorsal gill arch) 394–407; *Panturichthys fowleri* (leptocephalus unreported; fig. *P. isognathus*) 484–493; *Physalaemus barrioi* (description; chondrocranium; internal oral features) 507–518; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.: unknown)* 554–569; *Scinax (littoreus, perpusillus)* (tadpole diet; oral morphology) 683–689; *Xenopus laevis* (tadpole growth/development; artificial environment) 690–697; *Pleurodeles waltl* (newt prey choice) 756–761.
- LATERAL LINE**, *Panturichthys fowleri* (absence of pores) 484–493.
- LIFE HISTORY**, *Incilius nebulifer* (larvae, salinity effects) 106–114; *Bitis schneideri* (seasonal movements, sexual/ontogenetic) 115–120; *Liolaemus poecilochromus* (ontogenetic diet shifts) 203–210; *Chaetodon (auriga, aureofasciatus, rainfordi, plebius)*, *Chelmon rostratus* (larval settlement behavior) 211–221; *Etheostoma in-scriptum* (reproduction; fecundity; sex ratio; GSI) 237–242; *Careproctus (lycopersicus n. sp., kamikawai n. sp., ovigerus)* (eggs; reproduction maturity) 257–265; *Etheostoma (forbesi, crossopterum)* (reproduction) 432–439; *Lithobates areolatus* (migration; home range; burrow philopatry; mortality/survivorship) 440–450; *Fundulus jenkinsi* (spawning season; oogenesis) 451–459; *Physalaemus barrioi* (advertisement call; tadpole morphology; calling/egg-laying sites) 507–518; *Physalaemus pustulosus* (acoustic environment; evoked vocal responses) 678–682; *Scinax (littoreus, perpusillus)* (tadpole diet) 683–689; *Percina nigrofasciata* (reproduction; population variation; clutch/egg size; seasonality) 714–721; *Lepidochelys olivacea* (hatchling emergence) 738–747.
- MERISTICS/MORPHOMETRICS**, *Plesiopoma (curvidens n. gen., n. sp.)* 6–11; *Tricheilostoma* (13 spp.) 37–48; *Loricaria (coximensis n. sp.)* (comparison w/ geographic. proximate spp.) 49–56; *Amolops (marmoratus, afghanus, indoburmanensis n. sp.)* 57–76; *Cathorops (nuchalis, spixii, wayuu n. sp.)* 77–97; *Clenocheirodon (pristis n. gen., n. sp.)* (male/female) 243–250; *Derilissus (lombardii n. sp.)* 251–256; *Careproctus (lycopersicus n. sp., kamikawai n. sp., ovigerus)* 257–265; *Hisonotus (bockmanni n. sp.)* 266–275; *Apterionotus (lindalvae n. sp.)* 283–292; *Awanois guamensis* (Guam/Hawaii/previous study) 293–300; *Pseudocrenilabrus multicolor* (gill arch, brooders/non-brooders/males) 382–388; *Hippocampus erectus* (morphometry & sexual dimorphism; weight, length) 389–393; *Hypostomus (careopinnatus n. sp.)* 494–500; *Astyanacinus (yariguies n. sp.)* (scaling analysis) 501–506; *Pareiorhaphis (proskynita n. sp.)* 519–526; *Adontosternarchus (duartei n. sp.)* 535–540; *Trinectes (hubbsbollinger n. sp.)* 541–546; *Peckoltia (feldbergae n. sp.)* 547–553; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.)* 554–569; *Congoglanis (howesi n. sp.)* 626–630; *Trichomycterus (minus n. sp.)* 631–636; *Nannoplecostomus (eleonora n. gen., n. sp.)* 637–647; *Guibemantis (tasifotsy n. sp., pulcher, liber)* 648–662; *Potamorrhaphis (labiatus n. sp.)* (canonical comp. analysis) 663–669; *Scoloplax (bailei n. sp.)* 670–677; *Scinax (littoreus, perpusillus)* (oral cavity comparison) 683–689.
- MESOPELAGIC**, Stomiidae (osteology; dorsal gill arch; phylogeny) 394–407.
- METHODS**, *Bitis schneideri* (seasonal movement; radio-telemetry) 115–120; *Mycteroperca venenosa* (sound record; remote video) 135–144; *Cryptobranchus alleganiensis bishopi* (radio-telemetry; captive-rearing) 150–159; *Clemmys guttata* (radio-telemetry; flood effects & movement/home range) 179–190; *Anaxyrus fowleri*, *Incilius nebulifer* (museum & GIS data) 307–320; *Lithobates (pipiens, sylvaticus)*, *Pseudacris maculata*, *Anaxyrus (cognatus, woodhousii)* (habitat suitability modeling) 321–330; *Ambystoma macrodactylum* (under-road tunnels; conservation) 331–340; *Alligator mississippiensis* (gastric lavage, hose-Heimlich; digestion rate) 419–423; *Lithobates areolatus* (telemetry; pre-/post-breeding movements) 440–450; *Lampropeltis g. getula* (radio-telemetry; diel activity) 460–464; *Scinax (alter sp. complex)* (morphology, canonical discriminant analysis) 554–569; *Ambystoma annulatum* (microsatellite DNA; polymorphic loci) 570–577; *Uta stansburiana* (population crosses; reproductive isolating mechanisms; speciation) 593–602; *Hollandichthys*, *Rachoviscus* (histology, spermiogenesis) 609–625; *Pituophis melanoleucus mugitus* (radio-telemetry; home range) 706–713; *Bitis schneideri* (radio-telemetry; movement patterns) 732–737.
- MICROSATELLITE DNA**, *Egernia striolata* (aggregation kinships) 130–134; *Ambystoma annulatum* (polymorphic loci) 570–577.
- MIGRATION**, *Ambystoma macrodactylum* (breeding sites; precipitation; under-road tunnel usage; mortality) 331–340; *Lithobates sylvaticus* (juvenile orientation; olfactory cues) 424–431; *Lithobates areolatus* (pre-/post-breeding; telemetry; burrow philopatry) 440–450.
- MINIATURIZATION**, *Nannoplecostomus (eleonora n. gen., n. sp.)* (smallest known loricariid; summary list) 637–647.
- MITOCHONDRIAL DNA**, *Rypticus (carpenteri n. sp., subiffrenatus)* (distinction; consensus tree) 23–36; *Amolops (marmoratus, afghanus, panhai, indoburmanensis n. sp.)* (cryptic spp.) 57–76; *Hydrophis parviceps* (generic placement) 276–282; *Awanois (guamensis, stamineus, ocellaris)* (evolutionary lineages; phylogenetic analysis) 293–300; *Graptomys (o. ouachitensis, o. sabinensis, pseudogeographica)* (variation; haplotypes) 301–306; *Guibemantis (tasifotsy n. sp., pulcher, liber)* (Bayesian phylogeny) 648–662.
- MORPHOLOGY**, *Plesiopoma (curvidens n. gen., n. sp.)* (diagnosis; description; osteology; b&w photos., figs.; sexual dimorphism; coloration; morphometrics) 6–11; *Phymaturus (extrilidus n. sp.)* (diagnosis; holotype description; variation; coloration; b&w photos., color fig.) 12–22; *Rypticus (carpenteri n. sp., subiffrenatus)* (diagnosis; description; coloration; meristics; color photos.) 23–36; *Tricheilostoma (dimidiatum n. comb.)* (diagnosis;

- neotype description; morphometrics; variation; coloration; b&w photos.; head & hemipenis) 37–48; *Loricaria (coximensis n. sp.)* (diagnosis; description; b&w photo.; abdominal plates; sexual dimorphism; coloration) 49–56; *Amolops (marmoratus, afghanus, indoburmanensis n. sp.)* (diagnosis; description; coloration; b&w photos.; morphometrics) 57–76; *Cathorops (nuchalis, spixii, wayuu n. sp.)* (diagnosis; description; coloration; sexual dimorphism; b&w photos.; tooth plates), (meristics; PC analysis plots; key to east. So. Amer. spp.) 77–97; *Scarus ferrugineus* (ovaries/oogenesis, ultrastructure; b&w photos.) 229–236; *Ctenocheirodon* (n. gen.; diagnosis), (*pristis n. sp.*: diagnosis; description; coloration; sexual dimorphism; osteology; dentition; b&w/color photos.) 243–250; *Derilissus (lombardii n. sp.)* (diagnosis; description; meristics; color photos.; sucking disc papillae), (congener comparisons) 251–256; *Careproctus (hyopercus n. sp., kamikawai n. sp., ovigerus)* (diagnosis; description; osteology; meristics/morphometrics; color photos.) 257–265; *Hisonotus (bockmanni n. sp.)* (diagnosis; description; osteology; coloration; meristics; sexual dimorphism; b&w/color photos.) 266–275; *Hydrophis parviceps* (diagnosis; redescription; dentition; squamation; b&w/color photos.) 276–282; *Apterotonotus (lindavae n. sp.)* (diagnosis; description; coloration; sexual dimorphism; b&w photos., male/female) 283–292; *Atherinops affinis*, *Menidia menidia* (vertebral number, Jordan's Rule, latitudinal variation) 341–350; *Pseudocrenilabrus multicolor* (gill arch, mouth brooders/non-brooders; meristics; photo.) 382–388; *Hippocampus erectus* (sexual dimorphism; allometry & morphometry) 389–393; Stomiidae (dorsal gill arches; ontogeny; synapomorphy; color photos., figs.) 394–407; *Bathyaethiops (atercrinis n. sp.)* (diagnosis; description; coloration; osteology; meristics/morphometrics; color photos.) 478–483; *Panturichthys fowleri* (redescription; meristics/morphometrics; viscera; dentition; osteology; variation; b&w photos./figs.), Heterenchelyidae (key to spp.) 484–493; *Hypostomus (careopinnatus n. sp.)* (diagnosis; description; coloration; b&w photos.) 494–500; *Astyanacinus (yariguies n. sp.)* (diagnosis; description; coloration; dentition; morphometrics; b&w photo.) 501–506; *Physalaemus barrooi* (redescription; coloration; karotype; tadpole; osteology) 507–518; *Pareiorhaphis (proskynita n. sp.)* (diagnosis; description; coloration; sexual dimorphism; meristics/morphometrics; color photo.) 519–526; *Trypauchenopsis intermedia* (gen. & sp. redescription; coloration; osteology; sex dimorphism; head processes; figs.) 527–534; *Adontosternarchus* (key to spp.), (*duartei n. sp.*) (diagnosis; description; coloration; b&w photo.) 535–540; *Trinectes (hubbsbollinger n. sp.)* (diagnosis; description; coloration; morphometrics; fimbria; color photos.) 541–546; *Peckoltia (feldbergae n. sp.)* (diagnosis; description; coloration; morphometrics; sexual dimorphism; variation; b&w/color photos.) 547–553; *Scinax (alter)* (holotype redescription), (*imbegue n. sp., tymbamirim n. sp.*) (diagnosis; describes.), (coloration; variation; tadpole) 554–569; *Uta stansburiana* (polymorphism; speciation) 593–602; †*Vidalamini* (new fossil specimen; description; maxilla; b&w photo./figs.) 603–608; *Hollandichthys*, *Rachoviscus* (sperm) 609–625; *Congoglanis (howesi n. sp.)* (diagnosis; description; coloration; morphometrics; photo.), (key to genus) 626–630; *Trichomycterus (minus n. sp.)* (diagnosis; description; coloration; morphometrics; photo) 631–636; *Nannoplecostomus (eleonora n. gen., n. sp.)* (diagnosis; description; coloration; miniaturization; osteology; sexual dimorphism; photo./figs.) 637–647; *Guibemantis (tasifotsy n. sp.)* (diagnosis; description; coloration; morphometrics; photos.) 648–662; *Potamorrhaphis (labiatus n. sp.)* (diagnosis; description; coloration; sexual dimorphism; photos./figs.), (key to spp.) 663–669; *Scoloplax (baileyi n. sp.)* (diagnosis; description; coloration; morphometrics; osteology; dentition; photos./figs.), (key to spp.) 670–677; *Scinax (littoreus, perpusillus)* (tadpole oral morphology; photo.) 683–689; *Plethodon cinereus* (eco-morph color change) 748–755.
- MOVEMENT.** *Bitis schneideri* (seasonal, sexual/ontogenetic) 115–120; *Clemmys guttata* (patterns, pre-/post-flooding) 179–190; *Acris crepitans* (effect of habitat type) 191–196; *Chaetodon (auriga, aureofasciatus, rainfordi, plebius)*, *Chelmon rostratus* (larval swimming speeds) 211–221; *Ambystoma macrodactylum* (breeding site migrations; conservation, under-road tunnels) 331–340; *Lithobates areolatus* (burrow philopatry; telemetry) 440–450; *Basiliscus plumifrons* (activity budget; habitat usage; sexual/ontogenetic) 465–471; *Pituophis melanoleucus mugitus* (home range; refugia; road crossing; seasonality) 706–713; *Bitis schneideri* (patterns; seasonality) 732–737.
- MUSEUMS.** *Crotalus cerastes* (feeding patterns & reproduction) 100–105; *Graptemys (o. ouachitensis, o. sabinensis, pseudogeographica)* (tissue samples; mtDNA analysis) 301–306; *Anaxyrus fowleri*, *Incilius nebulifer* (distributions; historical data) 307–320.
- NATURAL HISTORY.** *Phymaturus (extrilidus n. sp.)* (basking & body temperatures; parental care; female/juvenile associations) 12–22; *Crotalus cerastes* (seasonal aphasia; reproduction) 100–105; *Incilius nebulifer* (salinity levels & hatching/metamorphosis, survival) 106–114; *Bitis schneideri* (seasonal movement) 115–120; *Eleutherodactylus planirostris* (introduced sp., diet, native/non-native invert. spp.) 121–129; *Egernia striolata* (aggregations; juvenile dispersal) 130–134; *Mycteroperca venenosa* (social/reproduction behavior; color phases) 135–144; *Sibon (longifrenis, argus, annulatus, nebulatus)*, *Dipsas (nicholsi, sp.)* (diet) 197–202; *Liolaemus poecilochromus* (diet; ontogeny; variation) 203–210; *Chaetodon (auriga, aureofasciatus, rainfordi, plebius)*, *Chelmon rostratus* (ontogeny; larval settlement/habitat selection; movement) 211–221; *Hydrophis parviceps* (reproduction; feeding; counter-coloration) 276–282; *Lampropeltis g. getula* (diel activity) 460–464; *Peckoltia (feldbergae n. sp.)* (rheophilic; nocturnal foraging; diet) 547–553; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.)* (habitat; breeding season; diet; fecundity; calls) 554–569; *Uta stansburiana* (reproductive isolating mechanism btwn. populations; mating strategies) 593–602; *Guibemantis (tasifotsy n. sp., pulcher, liber)* (breeding sites; calls) 648–662; *Potamorrhaphis (labiatus n. sp.)* (ontogeny) 663–669; *Pituophis melanoleucus mugitus* (home range; movement; variation; habitat selection; refugia) 706–713; *Bitis schneideri* (movement patterns) 732–737.
- NESTS/NESTING.** *Clemmys guttata* (site selection, pre-/post-flooding) 179–190; *Gopherus agassizii* (anthropogenic chgs.; clutch size; hatching success) 222–228; *Etheostoma (forbesi, crossopterus)* (activity; clutch size; chronology) 432–439; *Lepidocheilus olivacea* (hatchling success; sand temperature; sex ratio) 738–747.
- NOMENCLATURE.** *Plesiopoma (curvidens n. gen., n. sp.)* 6–11; *Phymaturus (extrilidus n. sp.)* 12–22; *Rhyticus (carpenteri n. sp.)* 23–36; *Tricheilostoma (dimidiatum n. comb.)* (neotype design.), (*jani n. sp.*) 37–48; *Loricaria (coximensis n. sp.)* 49–56; *Amolops (marmoratus: neotype design.), (afghanus: lectotype design.), (indoburmanensis n. sp.)* 57–76; *Cathorops* (east. So. Amer. spp. synonymy; summary list), (*wayuu n. sp.*) 77–97; *Triakis semifasciata* (sr. syn.; *Mustelus felis* status) 98–99; *Ctenocheirodon (pristis n. gen., n. sp.)* 243–250; *Derilissus (lombardii n. sp.)* 251–256; *Careproctus (hyopercus n. sp., kamikawai n. sp.)* 257–265; *Hisonotus (bockmanni n. sp.)* 266–275; *Apterotonotus (lindavae n. sp.)* 283–292; *Awaous stamineus* (sp. status) 293–300; *Bathyaethiops (atercrinis n. sp.)* 478–483; *Panturichthys fowleri* (holotype status), Heterenchelyidae (genera/species w/ synonyms) 484–493; *Hypostomus (careopinnatus n. sp.)* 494–500; *Astyanacinus (yariguies n. sp.)* 501–506; *Pareiorhaphis (proskynita n. sp.)* 519–526; *Trypauchenopsis* (resurrection; spp. synonyms) 527–534; *Adontosternarchus (duartei n. sp.)* 535–540; *Trinectes (hubbsbollinger n. sp.)* 541–546; *Peckoltia (feldbergae n. sp.)* 547–553; *Scinax (alter, imbegue n. sp., tymbamirim n. sp.)* 554–569; *Congoglanis (howesi n. sp.)* 626–630; *Trichomycterus (minus n. sp.)* 631–636; *Nannoplecostomus (eleonora n. gen., n. sp.)* (Loricariidae, *incertae sedis*) 637–647; *Guibemantis (tasifotsy n. sp.)* 648–662; *Potamorrhaphis (labiatus n. sp.)* 663–669; *Scoloplax (baileyi n. sp.)* 670–677.
- NON-INDIGENOUS SPECIES.** *Eleutherodactylus planirostris* (Hawaii) 121–129; *Incilius nebulifer* (Louisiana) 307–320.
- OBITUARIES.** James Edward Craddock 160–161; Joseph T. Collins 351–354; John Michael Fitzsimons 762–764.
- OLFACTION.** *Gekko japonicus* (non-volatile chem. cues) 145–149; *Lithobates sylvaticus* (orientation cues) 424–431; Boidae, Colubridae (tongue-licking, chemoreception; intraspecific recognition) 472–477.

- ONTOGENY**, *Liolaemus poecilochromus* (diet, variation wrt growth) 203–210; *Chaetodon (aureofasciatus, rainfordi, auriga)* (larval settlement/swimming speeds) 211–221; Stomiidae (dorsal gill arch; morphology; synapomorphy) 394–407; *Basiliscus plumifrons* (activity budget) 465–471; *Potamorrhaphis (labiatus n. sp.)* (juvenile/adult morphology) 663–669.
- OOGENESIS**, *Scarus ferrugineus* (histology; ultrastructure; b&w photos.) 229–236; *Fundulus jenkinsi* (histology; color photos.) 451–459.
- ORIENTATION**, *Lithobates sylvaticus* (olfactory cues, juveniles) 424–431; *Lithobates areolatus* (burrow philopatry; telemetry; home range/movement) 440–450.
- OSTEOLOGY**, *Plesiopoma (curvidens n. gen., n. sp.)* (pelvic girdle; premax. tooth series) 6–11; *Cathorops (nuchalis, spixii, wayuu n. sp.)* (male/female tooth plates) 77–97; *Ctenocheirodon (pristis n. gen., n. sp.)* (procurent C rays, A rays; b&w photos.) 243–250; *Hisonotus (bockmanni n. sp.)* (skull, suspensorium, pelvic girdle) 266–275; *Atherinops affinis*, *Menidia menidia* (vert. number; Jordan's Rule) 341–350; Stomiidae (dorsal gill arches; ontogeny) 394–407; *Bathyaethiops (atercrinis n. sp.)* (teeth, anal fin, vertebrae), (comparison w/ *Brachypetersius* spp.) 478–483; *Panturichthys fowleri* (descriptions; variations) 484–493; *Physalampus barroii* (tadpole chondrocranium) 507–518; *Trypauchenopsis intermedia* (vertebrae, pterygiophores) 527–534; †*Vidalamiini* (maxilla; figs.) 603–608; *Nannoplecostomus (eleonora n. gen., n. sp.)* (head, pectoral girdle), Hypostominae (diagnosis characters) 637–647; *Scoloplax (baileyi n. sp.)* (unique tooth morphology; body plates; neurocranium) 670–677.
- PAEDOMORPHISM**, *Loricaria (coximensis n. sp.)* (sm. size at sexual maturity; incomplete/fewer plates) 49–56.
- PALEONTOLOGY**, †*Vidalamiini* (Eocene; osteology; habitat) 603–608.
- PARASITES**, *Liolaemus poecilochromus* (gut nematodes) 203–210.
- PARENTAL CARE**, *Phymaturus (extrilidus n. sp.)* (female/juvenile association) 12–22; *Pseudocrenilabrus multicolor* (mouth brooder; gill morphology) 382–388; *Hyalinobatrachium aureoguttatum* (male clutch attendance) 722–731.
- PATHOLOGY**, *Cryptobranchius alleganiensis bishopi* (captive-reared survival in wild; body condition) 150–159.
- PHYLOGENETIC ANALYSIS**, *Amolops (marmoratus afghanus, panhai, indoburmanensis + 17 spp.)* (cryptic sp. diversity; mtDNA, morphology; PC analysis; consensus tree) 57–76; *Rhycolius (carpenteri n. sp., subbifrenatus, saponaceus)* (mtDNA analysis tree; morphology) 23–36; *Ctenocheirodon* (n. gen.) (w/in Cheirodontinae & w/in Cheirodontini) 243–250; *Awaous (guamensis, stamineus)* (mtDNA, morphology; evolutionary lineages; taxonomic status) 293–300; *Graptemys (o. ouachitensis, o. sabinensis, pseudogeographica)* (mtDNA; variation; haplotypes) 301–306; *Guibemantis (tasifotsy n. sp., pulcher, liber)* (mtDNA; morphometrics) 648–662.
- PHYLOGENY**, Stomiidae (dorsal gill arches; morphology, homology) 394–407.
- PHYSIOLOGY**, *Phymaturus (extrilidus n. sp.)* (diurnal body temp.) 12–22; *Crotalus cerastes* (seasonal aphasia; reproduction) 100–105; *Incilius nebulifer* (effects salinity on early life stages) 106–114; *Clemmys guttata* (aestivation) 179–190; *Scarus ferrugineus* (oogenesis) 229–236; *Etheostoma inscriptum* (reproduction; GSI) 237–242; *Alligator mississippiensis* (digestion rate) 419–423; *Hollandichthys*, *Rachoviscus* (spermiogenesis) 609–625; *Graptemys flavimaculata* (stress response; corticosterone levels) 698–705.
- POPULATIONS**, *Egernia striolata* (tree-dwelling aggregations; kinship) 130–134; *Cryptobranchius alleganiensis bishopi* (survival released captive-reared populations; habitat; body condition) 150–159; *Graptemys (o. ouachitensis)* (Scioto R., OH; disjunct) 301–306; *Ambystoma macrodactylum* (Linnet Lake, Alb., Canada; movement patterns) 331–340; *Graptemys geographica*, *Trachemys scripta elegans* (habitat degradation, population decline) 367–374; *Uta stansburiana* (reproductive isolating mechanisms; speciation) 593–602; *Percina nigrofasciata* (population comparisons; reproductive variation) 714–721.
- PREDATION**, *Eleutherodactylus planirostris* (introduced; diet, native/non-native inverts.) 121–129; *Graptemys geographica* (river otter re-introduction) 367–374; *Pleurodeles walli* (newt prey choice) 756–761.
- REPRODUCTION**, *Crotalus cerastes* (seasonal aphasia) 100–105; *Mycteroperca venenosa* (spawning behavior; color phases; sound production) 135–144; *Gopherus agassizii* (nesting; clutch size; hatching success) 222–228; *Scarus ferrugineus* (oogenesis; ultrastructure; b&w photos.) 229–236; *Etheostoma inscriptum* (fecundity; GSI; sex ratio) 237–242; *Hydrophis parviceps* (embryos; parturition) 276–282; *Pseudocrenilabrus multicolor* (mouth brooding & gill morphology) 382–388; *Hippocampus erectus* (sexual dimorphism & selection) 389–393; *Etheostoma (forbesi, crossopterus)* (nesting; clutch size; hybridization) 432–439; *Fundulus jenkinsi* (oogenesis; GSI; spawning season) 451–459; *Scinax (tybamirim n. sp.)* (breeding season; clutch size) 554–569; *Uta stansburiana* (reproductive isolating mechanism, speciation) 593–602; *Hollandichthys*, *Rachoviscus* (spermiogenesis) 609–625; *Guibemantis (tasifotsy n. sp., pulcher, liber)* (breeding site selection) 648–662; *Percina nigrofasciata* (population variation; seasonality; clutch/egg size) 714–721; *Hyalinobatrachium aureoguttatum* (microhabitat selection; male clutch attendance) 722–731.
- RESPIRATION**, *Pseudocrenilabrus multicolor* (mouth brooding & gill morphology) 382–388; *Xenopus laevis* (tadpole growth/development; artificial environment) 690–697.
- SALINITY**, *Incilius nebulifer* (effect on early life stages), (summary table of studies) 106–114.
- SENSORY**, *Lithobates sylvaticus* (olfactory cues, orientation) 424–431; Boidae, Colubridae (chemoreception, intraspecific recognition) 472–477.
- SETTLEMENT**, *Chaetodon (auriga, aureofasciatus, rainfordi, plebius)*, *Chelmon rostratus* (habitat selection) 211–221.
- SEX**, *Liolaemus poecilochromus* (diet variation, males/females) 203–210; *Atherinops affinis*, *Menidia menidia* (determination, TSD vs. GSD; latitudinal variation) 341–350.
- SEX RATIO**, *Clemmys guttata* (pre-/post-flooding) 179–190; *Etheostoma inscriptum* (females favored) 237–242; *Atherinops affinis* (effects of temperature/latitude change) 341–350; *Graptemys geographica* (btwn.-year comparison) 367–374; *Lepidocheilus olivacea* (hatchling success; sand temperature) 738–747.
- SEXUAL DIMORPHISM**, *Plesiopoma (curvidens n. gen., n. sp.)* 6–11; *Phymaturus (extrilidus n. sp.)* (not strong) 12–22; *Tricheilostoma (jani n. sp., dimidiatum n. comb.)* (scales, total length) 37–48; *Loricaria (coximensis n. sp.)* (papillae; pectoral spine, tooth cusps) 49–56; *Amolops (marmoratus, afghanus, indoburmanensis n. sp.)* (size; PCA values) 57–76; *Cathorops (nuchalis, spixii, wayuu n. sp.)* (size, tooth plates) 77–97; *Mycteroperca venenosa* (dichromatic) 135–144; *Liolaemus poecilochromus* (males larger) 203–210; *Etheostoma inscriptum* (males larger) 237–242; *Ctenocheirodon (pristis n. gen., n. sp.)* (A fin lepidotrichia, ventral procurent C rays) 243–250; *Hisonotus (bockmanni n. sp.)* (male urogenital papillae) 266–275; *Hydrophis parviceps* (scales, vertebrae, position of heart) 276–282; *Apteronotus (lindalvae n. sp.)* (males larger; PC analysis) 283–292; *Hippocampus erectus* (allometry & morphometry) 389–393; *Pareiorhaphis (proskynita n. sp.)* (male odontodes, fleshy head lobes) 519–526; *Trypauchenopsis intermedia* (male genital papillae) 527–534; *Peckoltia (feldbergae n. sp.)* (male odontodes) 547–553; *Nannoplecostomus (eleonora n. gen., n. sp.)* (urogenital opening; pelvic-fin flap) 637–647; *Potamorrhaphis (labiatus n. sp.)* (genital papilla) 663–669.
- SIZE**, *Loricaria (coximensis n. sp.)* (paedomorphic) 49–56; *Amolops (marmoratus, afghanus, indoburmanensis n. sp.)* (sexual dimorphism) 57–76; *Liolaemus poecilochromus* (smallest herbivorous reptile) 203–210; *Graptemys geographica* (decline in large adult females) 367–374; *Trypauchenopsis intermedia* (male/female freq. distrib.) 527–534; *Nannoplecostomus (eleonora n. gen., n. sp.)* (smallest loriciariid; summary list) 637–647.
- SOUND**, *Mycteroperca venenosa* (social/reproduction behavior; oscillo-/spectrograms) 135–144; *Gekko japonicus* (acoustic behavior; intra-/intersexual spectrograms) 145–149.
- SPECIATION**, *Uta stansburiana* (reproductive isolating mechanism) 593–602.

SPERMIOGENESIS, *Hollandichthys*, *Rachoviscus* (sperm competitiveness) 609–625.

SURVIVAL, *Incilius nebulifer* (effects of salinity levels on early life stages) 106–114; *Cryptobranchus alleganiensis bishopi* (captive-reared/released; body condition) 150–159; *Lithobates areolatus* (movement; mortality/survivorship) 440–450; *Lepidochelys olivacea* (hatchling emergence success, *in situ* & relocated nests) 738–747.

SYSTEMATICS, *Plesiopotopoma* (*curvidens* n. gen., n. sp.) (placement w/in Hypoptopomatinae; comparison w/ Neoplecostominae) 6–11; *Phymaturus* (*extrilidus* n. sp.) (placement w/in *palluma* gr.) 12–22; *Rhyticus* (*carpenteri* n. sp.) (separated from *subbifrenatus*) 23–36; *Tricheilostoma* (*jani* n. sp., *dimidiatum* n. comb.) 37–48; *Loricaria* (*coximensis* n. sp.) (comparison w/ geographic proximate spp.) 49–56; *Amolops* (Myanmar, *marmoratus* sp. complex; phylogenetic analysis) 57–76; *Cathorops* (east. So. Amer. spp.) (PC analysis, synonymy; key to spp.) 77–97; *Clenocheirodon* (*pristis* n. gen., n. sp.) (w/in Cheirodontinae & w/in Cheirodontini) 243–250; *Derilissus* (*lombardii* n. sp.) (apomorphic character) 251–256; *Careproctus* (*lycopersicus* n. sp., *kamikawai* n. sp., *ovigerus*) (comparisons) 257–265; *Hisonotus* (intrageneric comparisons) 266–275; *Apteranotus* (*lindalvae* n. sp.) (generic placement) 283–292; *Awaous* (*guamensis*, *stamineus*) (evolutionary lineages) 293–300; *Bathyaethiops* (*atercrinis* n. sp.) (comparison w/ *Brachypetersius*) 478–483; *Panturichthys fowleri* (key to family; synonymy) 484–493; *Hypostomus* (*careopinatus* n. sp.) (placement w/in genus) 494–500; *Asryanacinus* (generic diagnosis characters) 501–506; *Pareiorhaphis* (*proskynita* n. sp.) (congener comparisons) 519–526; *Trypauchenopsis intermedia* (placement w/in “*Taenioides*” group) 527–534; *Adontosternarchus* (key to spp.) 535–540; *Peckoltia* (*feldbergae* n. sp.) (generic status wrt *Hemiancistrus*) 547–553; *Scinax* (*alter*, *imbegue* n. sp., *tybamirim* n. sp.) (species complex) 554–569; *Congoglanis* (key to spp.) 626–630; *Trichomycterus* (*minus* n. sp.) (comparison to congeners) 631–636; *Nannoplecostomus* (*eleonorae* n. gen., n. sp.) (loricariid, *incertae sedis*) 637–647; *Guibemantis* (*tasifotsy* n. sp., *pulcher*, *liber*) (phylogenetic analysis; mtDNA; morphometrics) 648–662; *Potamorhaphis* (key to spp.) 663–669; *Scoloplax* (key to spp.) 670–677.

TAXONOMY, *Plesiopotopoma* (*curvidens* n. gen., n. sp.) 6–11; *Phymaturus* (*extrilidus* n. sp.) (comparison w/in *palluma* gr.) 12–22; *Rhyticus* (*carpenteri* n. sp.) (separate from *subbifrenatus*) 23–36; *Tricheilostoma* (*jani* n. sp., *dimidiatum* n. comb.) (comparison w/ Neotropical leptotyphlopids) 37–48; *Loricaria* (*coximensis* n. sp.) 49–56; *Amolops* (*marmoratus*: description neotype), (*afghanus*: description lectotype), (*indoburmanensis* n. sp.) 57–76; *Cathorops* (east. So. Amer. spp.; synonymy), (*wayuu* n. sp.) 77–97; *Triakis semifasciata* (senior synonym; *Mustelus felis* invalidated) 98–99; *Clenocheirodon* (n. gen.) (subfamily/tribe relationships) 243–250; *Derilissus* (*lombardii* n. sp.) (congener comparisons) 251–256; *Careproctus* (*lycopersicus* n. sp., *kamikawai* n. sp., *ovigerus*) (comparisons) 257–265; *Hisonotus* (*bockmanni* n. sp.) (congener comparison) 266–275; *Hydrophis parviceps* (redescription; morphology & mtDNA data) 276–282; *Apteranotus* (nominal spp., table), (*lindalvae* n. sp.)

(generic placement) 283–292; *Awaous* (*guamensis*, *stamineus*) (cryptic spp.; evolutionary lineages) 293–300; Stomiidae (dorsal gill arch, synapomorphy) 394–407; *Bathyaethiops* (*atercrinis* n. sp.) (osteological autapomorphies) 478–483; *Panturichthys fowleri* (redescription; key to Heterenchelyidae) 484–493; *Hypostomus* (*careopinatus* n. sp.) (generic synapomorphies) 494–500; *Asryanacinus* (generic characters) 501–506; *Physalasmus barrioi* (redescription, adult/tadpole) 507–518; *Pareiorhaphis* (*proskynita* n. sp.) (description; comparison to congeners & other loricariids) 519–526; *Trypauchenopsis intermedia* (redescription; relationships) 527–534; *Adontosternarchus* (*duartei* n. sp.) (description; key to spp.) 535–540; *Trinectes* (*hubbsboller* n. sp.) (description; color photos.) 541–546; *Peckoltia* (*feldbergae* n. sp.) (description; comparison w/ congeners) 547–553; *Scinax* (*alter*, *imbegue* n. sp., *tybamirim* n. sp.) (descrips.; comparison to congeners) 554–569; *Congoglanis* (*howesi* n. sp.) (description; key to spp.) 626–630; *Trichomycterus* (*minus* n. sp.) (description; comparison to congeners) 631–636; *Nannoplecostomus* (*eleonorae* n. gen., n. sp.) (description) 637–647; *Guibemantis* (*tasifotsy* n. sp.) (description) 648–662; *Potamorhaphis* (*labiatus* n. sp.) (description), (key to spp.) 663–669; *Scoloplax* (*baileyi* n. sp.) (description), (key to spp.) 670–677.

TEMPERATURE, *Atherinops affinis*, *Menidia menidia* (latitudinal variation; effect on vertebral number, sex determination) 341–350; *Bothrops insularis* (thermoregulation; seasonal variation; habitat usage) 408–418.

TROPHIC NICHE, *Scinax* (*littoreus*, *perpusillus*) (comparison) 683–689.

VARIATION, *Plesiopotopoma* (*curvidens* n. gen., n. sp.) (table, morphometrics; comparison, lg./sm. sizes) 6–11; *Phymaturus* (*extrilidus* n. sp.) (morphometrics; coloration, sexual/ontogenetic) 12–22; *Tricheilostoma* (*jani* n. sp., *dimidiatum* n. comb.) (female/male scales & lengths) 37–48; *Amolops* (*marmoratus*, *afghanus*, *indoburmanensis* n. sp.) (male/female morphometrics; tables) 57–76; *Egernia striolata* (factors influence aggregation behavior) 130–134; *Liolaemus poecilochromus* (diet; ontogenetic shift, sexual/ontogenetic) 203–210; *Graptemys o. ouachitensis* (mtDNA; haplotypes) 301–306; *Ambystoma macrodactylum* (movement to breeding sites; moisture; under-road tunnel usage) 331–340; *Atherinops affinis*, *Menidia menidia* (Jordan's Rule, latitudinal temperature; effects on vertebral number/sex determination) 341–350; *Bothrops insularis* (thermoregulation; seasonal variation & behavior) 408–418; *Basiliscus plumifrons* (activity budget: seasonal, habitat, ontogenetic) 465–471; *Panturichthys fowleri* (morphometrics; dentition) 484–493; *Peckoltia* (*feldbergae* n. sp.) (morphological) 547–553; *Scinax* (*alter*, *imbegue* n. sp., *tybamirim* n. sp.) (morphological; canonical discriminant analysis) 554–569; *Guibemantis* (*tasifotsy* n. sp., *pulcher*) (coloration) 648–662; *Graptemys flavimaculata* (cortisol levels; seasonality) 698–705; *Pituophis melanoleucus mugitus* (home range) 706–713; *Percina nigrofasciata* (reproduction; population variation) 714–721; *Bitis schneideri* (movement patterns, sexual/seasonal) 732–737.

VICARIANCE, *Graptemys o. ouachitensis* (Pleistocene glacial refugia) 301–306.

VOLUME CONTENTS

Copeia 2012, Nos. 1–4

Abdel-Aziz, Sayeda H., Mashael Al-Otaibi, Amany M. Osman, T. El-Sayed Ali, and Fayza Bawazeer. Ultrastructural Evidence of Oogenesis of the Rusty Parrotfish, <i>Scarus ferrugineus</i> (Teleostei: Scaridae)	229	Booth, David T. (see Maulany, Risma I.)	
Acero P., Arturo (see Marceniuk, Alexandre P.)		Bovo, Rafael P., Otavio A. V. Marques, and Denis V. Andrade. When Basking Is Not an Option: Thermoregulation of a Viperid Snake Endemic to a Small Island in the South Atlantic of Brazil	408
Adams, Dean C. (see Kraemer, Andrew C.)		Branco, Christina W. C. (see Sabagh, Leandro T.)	
Al-Otaibi, Mashael (see Abdel-Aziz, Sayeda H.)		Briggler, Jeffrey T. (see Bodinof, Catherine M.)	
Alexander, Graham J. (see Maritz, Bryan)		Britto, Marcelo R. (see Pereira, Edson H. L.)	
Alexander, Graham J. (see Maritz, Bryan)		Brodie, Bekka S. (see Popescu, Viorel D.)	
Alexander, Laura G., Simon P. Lailvaux, Joseph H. K. Pechmann, and Philip J. DeVries. Effects of Salinity on Early Life Stages of the Gulf Coast Toad, <i>Incilius nebulifer</i> (Anura: Bufonidae)	106	Brown-Peterson, Nancy J. (see Lang, Erik T.)	
Ali, T. El-Sayed (see Abdel-Aziz, Sayeda H.)		Brown, Ashley D., Kathleen Temple-Miller, Willem M. Roosenburg, and Matthew M. White. Mitochondrial DNA Variation in the Ouachita Map Turtle	301
Anderson, Paul A. Sexual Dimorphism in Morphometry and Allometry of the Adult Lined Seahorse, <i>Hippocampus erectus</i>	389	Burns, John R. (see Quagio-Grassiotto, Irani)	
Andrade, Denis V. (see Bovo, Rafael P.)		Calich, Hannah J., and Richard J. Wassersug. The Architecture of the Aquatic Environment and Its Influence on the Growth and Development of Tadpoles (<i>Xenopus laevis</i>)	690
Andreone, Franco (see Lehtinen, Richard M.)		Carvalho, Murilo, and Aléssio Datovo. A New Species of Cascudinho of the Genus <i>Hisonotus</i> (Siluriformes: Loricariidae: Hypoptopomatinae) from the Upper Rio Tapajós Basin, Brazil	266
Appeldoorn, Richard S. (see Schärer, Michelle T.)		Castro-Herrera, Fernando (see Valencia-Aguilar, Anyelet)	
Arias, Federico (see Valdecantos, María Soledad)		Cavallaro, Marcel R. (see Rodriguez, Mónica S.)	
Arundel, Terence R. (see Ennen, Joshua R.)		Chakrabarty, Prosanta, and Valerie Derouen. Fishes: The Animal Answer Guide. By Gene Helfman and Bruce Collette. Do Fish Sleep? Fascinating Answers to Questions About Fishes. By Judith S. Weis	361
Azevedo, Marco A. (see Quagio-Grassiotto, Irani)		Chakrabarty, Prosanta. The Biology of Gobies. By Robert A. Patzner, James L. Van Tassel, Marcelo Kovačić, and B. G. Kapoor	359
Baicere-Silva, Clarianna Martins (see Quagio-Grassiotto, Irani)		Chapleau, François (see Duplain, René R.)	
Baldwin, Carole C., and Lee A. Weigt. A New Species of Soapfish (Teleostei: Serranidae: <i>Rypticus</i>), with Redescription of <i>R. subbifrenatus</i> and Comments on the Use of DNA Barcoding in Systematic Studies	23	Chapman, Lauren J. (see O'Connor, Constance M.)	
Barichivich, William (see Nifong, James C.)		Combs, Daniel L. (see Bergen, Darrin R.)	
Baumann, Hannes, Jorge A. Rosales Casán, and David O. Conover. Contrasting Latitudinal Variations in Vertebral Number and Sex Determination in Pacific Versus Atlantic Silverside Fishes	341	Connette, Grant M. (see Peterman, William E.)	
Bawazeer, Fayza (see Abdel-Aziz, Sayeda H.)		Conover, David O. (see Baumann, Hannes)	
Baxter, Greg S. (see Maulany, Risma I.)		Corl, Ammon, Lesley T. Lancaster, and Barry Sinervo. Rapid Formation of Reproductive Isolation between Two Populations of Side-Blotched Lizards, <i>Uta stansburiana</i>	593
Beard, Karen H. (see Olson, Christina A.)		Cox Fernandes, Cristina (see de Santana, Carlos David)	
Bell, Michael A. Catalogue of the Neogene Bony Fishes of California. A Systematic Inventory of All Published Accounts. By Harry L. Fierstine, Richard W. Huddleston, and Gary T. Takeuchi	582	Datovo, Aléssio (see Carvalho, Murilo)	
Bergen, Darrin R., Daniel L. Combs, and Hayden T. Mattingly. Comparative Nesting Biology of Two Sympatric Darters (Percidae: <i>Etheostoma</i> : <i>Catnotus</i>) in Headwater Streams of the Caney Fork River, Tennessee	432	Dawson, Beth, and Michael J. Ryan. Evoked Vocal Responses Change with Experience in Male <i>Physalaemus pustulosus</i>	678
Beringer, Jeff (see Bodinof, Catherine M.)		de C. Rossa-Feres, Denise (see Provete, Diogo B.)	
Betancur-R., Ricardo (see Marceniuk, Alexandre P.)		de Oliveira, Renildo R., Lucia Rapp Py-Daniel, Jansen Zuanon, and Marcelo S. Rocha. A New Species of the Ornamental Catfish Genus <i>Peckoltia</i> (Siluriformes: Loricariidae) from Rio Xingu Basin, Brazilian Amazon	547
Bjurlin, Curtis (see Ennen, Joshua R.)		de Santana, Carlos David, and Cristina Cox Fernandes. A New Species of Sexually Dimorphic Electric Knifefish from the Amazon Basin, Brazil (Gymnotiformes: Apteronotidae)	283
Blum, Michael J. (see Lindstrom, Daniel P.)		de Santana, Carlos David, and Richard P. Vari. New Species of <i>Adontosternarchus</i> (Gymnotiformes, Apteronotidae) from the Rio Purus Basin, Brazil	535
Bodinof, Catherine M., Jeffrey T. Briggler, Randall E. Junge, Tony Mong, Jeff Beringer, Mark D. Wanner, Chawna D. Schuette, Jeff Ettling, and Joshua J. Millsbaugh. Survival and Body Condition of Captive-Reared Juvenile Ozark Hellbenders (<i>Cryptobranchus alleganiensis bishopi</i>) Following Translocation to the Wild	150		
Boone, Michelle (see Ramirez, Edward A.)			

- Delapieve, Maria Laura S. (see Sant'Anna, Vivianne B.)
- Derouen, Valerie (see Chakrabarty, Prosanta)
- Dever, Jennifer A., Allison M. Fuiten, Özlen Konu, and Jeffery A. Wilkinson. Cryptic Torrent Frogs of Myanmar: An Examination of the *Amolops marmoratus* Species Complex with the Resurrection of *Amolops afghanus* and the Identification of a New Species 57
- DeVries, Philip J. (see Alexander, Laura G.)
- Dias, Natacha Y. N. (see Sabagh, Leandro T.)
- Doan, Tiffany M. Perú: Yaguas-Cotuhé. Rapid Biological and Social Inventories Report 23. By N. Pitman, C. Vriesendorp, D. K. Moskovits, R. von May, D. Alvira, T. Wachter, D. F. Stotz, and A. del Campo (eds.) 769
- Dodd, C. Kenneth, Jr. Reptile Biodiversity. Standard Methods for Inventory and Monitoring. By R. W. McDiarmid, M. S. Foster, C. Guyer, J. W. Gibbons, and N. Chernoff (eds.) 589
- Duckett, Paul E., Margaret H. Morgan, and Adam J. Stow. Tree-Dwelling Populations of the Skink *Egernia striolata* Aggregate in Groups of Close Kin 130
- Duplain, René R., François Chapleau, and Thomas A. Munroe. A New Species of *Trinectes* (Pleuronectiformes: Achiridae) from the Upper Río San Juan and Río Condoto, Colombia 541
- Ebert, David A. The Sharks of North America. By Jose I. Castro 765
- Eggert, Lori S. (see Peterman, William E.)
- Elmberg, Johan (see Rasmussen, Arne R.)
- Ennen, Joshua R., Jeffrey E. Lovich, Katherin P. Meyer, Curtis Bjurlin, and Terence R. Arundel. Nesting Ecology of a Population of *Gopherus agassizii* at a Utility-Scale Wind Energy Facility in Southern California 222
- Eschmeyer, William N. (see Pietsch, Theodore W.)
- Espinoza, Robert E. (see Lobo, Fernando)
- Espinoza, Robert E. (see Valdecantos, María Soledad)
- Ettling, Jeff (see Bodinof, Catherine M.)
- Euliss, Ned H., Jr. (see Mushet, David M.)
- Fernandes, Ronaldo (see Pinto, Roberta R.)
- Fernández, Luis, and Richard P. Vari. New Species of *Trichomycterus* (Teleostei: Siluriformes) from the Andean Cordillera of Argentina and the Second Record of the Genus in Thermal Waters 631
- Ferraris, Carl J., Jr. (see Vari, Richard P.)
- Ferreira, Gustavo Luis (see Sabagh, Leandro T.)
- Foltz, Jeffrey W. (see Irwin, Stephan W.)
- Font, William F. (see Schoenfuss, Heiko L.)
- Franz, Richard (see Miller, Gabriel J.)
- Fuiten, Allison M. (see Dever, Jennifer A.)
- Gabirot, Marianne, Pablo Picerno, Jorge Valencia, Pilar Lopez, and José Martín. Species Recognition by Chemical Cues in Neotropical Snakes 472
- Gagne, Roderick B. (see Lindstrom, Daniel P.)
- García-de-Lomas, Juan, José María Torres, Ricardo Reques, and Ignacio Hernández. *Pleurodeles waltli* Newt Larvae Diet Regarding Available Prey: Choice or Mere Coincidence? 756
- Garey, Michel V. (see Provete, Diogo B.)
- Gilliam, James F. (see Lindstrom, Daniel P.)
- Glaudas, Xavier (see Webber, Michael M.)
- Glaw, Frank (see Lehtinen, Richard M.)
- Gravlund, Peter (see Rasmussen, Arne R.)
- Greenfield, David W. Shore Fishes of Easter Island. By John E. Randall and Alfredo Cea 165
- Gruber, David F. (see Sparks, John S.)
- Haddad, Célio F. B. (see Provete, Diogo B.)
- Hartel, Karsten E., and Pamela Polloni. James Edward Craddock (1937–2009) 160
- Heemeyer, Jennifer L., and Michael J. Lannoo. Breeding Migrations in Crawfish Frogs (*Lithobates areolatus*): Long-Distance Movements, Burrow Philopatry, and Mortality in a Near-Threatened Species 440
- Heins, David C. (see Hughey, Myra C.)
- Heithaus, Michael R. (see Nifong, James C.)
- Henderson, Robert W. Invasive Pythons in the United States: Ecology of an Introduced Predator. By M. E. Dorcas and J. D. Willson 358
- Hernández, Emilton (see Torres-Mejía, Mauricio)
- Hernández, Ignacio (see García-de-Lomas, Juan)
- Hill, Robert V. (see O'Leary, Maureen A.)
- Howze, Jennifer M., and Lora L. Smith. Factors Influencing Eastern Kingsnake Diel Activity 460
- Hughey, Myra C., David C. Heins, Howard L. Jelks, Bridget A. Ory, and Frank Jordan. Variation in Reproductive Life History Traits between Two Populations of Blackbanded Darters (*Percina nigrofasciata*) 714
- Hunter, Malcolm L. (see Popescu, Viorel D.)
- Inui, Yoko (see Jono, Teppei)
- Irmak, Erhan (see Smith, David G.)
- Irwin, Stephan W., and Jeffrey W. Foltz. Reproductive Characteristics of the Turquoise Darter (*Etheostoma inscriptum*) in the Upper Piedmont of South Carolina 237
- Jawor, Jodie M. (see Selman, Will)
- Jelks, Howard L. (see Hughey, Myra C.)
- Jerep, Fernando C. (see Malabarba, Luiz R.)
- Johnson, G. David (see Schnell, Nalani K.)
- Johnson, Nathan A. (see Nifong, James C.)
- Johnson, Steve A. (see Miller, Gabriel J.)
- Jono, Teppei, and Yoko Inui. Secret Calls from under the Eaves: Acoustic Behavior of the Japanese House Gecko, *Gekko japonicus* 145
- Jordan, Frank (see Hughey, Myra C.)
- Junge, Randall E. (see Bodinof, Catherine M.)
- Kats, Lee B. Encyclopedia of Biological Invasions. By Daniel Simberloff and Marcel Rejmánek 169
- Kells, Val. Certainly More Than You Want to Know About The Fishes of The Pacific Coast—A Postmodern Experience. By Milton S. Love 360
- Kime, Nicole M. Animal Behavior. By Michael D. Breed and Janice Moore 774
- Kissner, Jessica (see Kraemer, Andrew C.)
- Konu, Özlen (see Dever, Jennifer A.)
- Kraemer, Andrew C., Jessica Kissner, and Dean C. Adams. Morphological Color-Change in the Red-Backed Salamander (*Plethodon cinereus*) While Kept in Captivity 748
- Kwet, Axel (see Nunes, Ivan)
- LaDuke, Thomas C. (see Lattanzio, Matthew S.)
- Lailvaux, Simon P. (see Alexander, Laura G.)
- Lancaster, Lesley T. (see Corl, Ammon)
- Lang, Erik T., Nancy J. Brown-Peterson, Mark S. Peterson, and William T. Slack. Seasonal and Tidally Driven Reproductive Patterns in the Saltmarsh Topminnow, *Fundulus jenkinsi* 451
- Langeani, Francisco (see Martins, Fernanda O.)
- Lannoo, Michael J. (see Heemeyer, Jennifer L.)
- Lattanzio, Matthew S., and Thomas C. LaDuke. Habitat Use and Activity Budgets of Emerald Basilisks (*Basiliscus plumifrons*) in Northeast Costa Rica 465

Lazzarotto, Henrique (see Rocha, Marcelo)	
Lehmann A., Pablo (see Reis, Roberto E.)	
Lehtinen, Richard M., Frank Glaw, Franco Andreone, Maciej Pabijan, and Miguel Vences. A New Species of Putatively Pond Breeding Frog of the Genus <i>Guibemantis</i> from Southeastern Madagascar	648
Leis, Jeffrey M., and Michelle N. Yerman. Behavior of Larval Butterflyfishes (Teleostei: Chaetodontidae) at Settlement on Coral Reefs	211
Lepre, Chris (see O'Leary, Maureen A.)	
Lima, Albertina P. (see Ribeiro, José W., Jr.)	
Lima, Flávio C. T. (see Ribeiro, Alexandre C.)	
Lindstrom, Daniel P., Michael J. Blum, Ryan P. Walter, Roderick B. Gagne, and James F. Gilliam. Molecular and Morphological Evidence of Distinct Evolutionary Lineages of <i>Awaous guamensis</i> in Hawai'i and Guam	293
Lips, Karen R. (see Ray, Julie M.)	
Litzgus, Jacqueline D. (see Yagi, Katharine T.)	
Lobo, Fernando, Robert E. Espinoza, Eduardo A. Sanabria, and Lorena B. Quiroga. A New <i>Phymaturus</i> (Iguania: Liolaemidae) from the Southern Extreme of the Argentine Puna	12
Locascio, James (see Schärer, Michelle T.)	
Lopez, Pilar (see Gabirot, Marianne)	
Lourenço, Luciana B. (see Provete, Diogo B.)	
Love, Milton S. The Curious Death of Peter Artedi. By Theodore W. Pietsch	162
Lovich, Jeffrey E. (see Ennen, Joshua R.)	
Magnusson, William E. (see Ribeiro, José W., Jr.)	
Mahon, Heidi K. (see Ray, Julie M.)	
Malabarba, Luiz R. (see Quaggio-Grassiotto, Irani)	
Malabarba, Luiz R., and Fernando C. Jerep. A New Genus and Species of Cheirodontine Fish from South America (Teleostei: Characidae)	243
Malhotra, Anita. Handbook of Venoms and Toxins of Reptiles. By Stephen P. Mackessy (ed.)	355
Malou, Raymond (see O'Leary, Maureen A.)	
Manonekene, Victor, and Melanie L. J. Stiassny. A New <i>Bathyaethiops</i> (Characiformes: Alestidae) from the Lékoumou River (Kouilou-Niari Basin) in the Republic of Congo; First Record of the Genus in the Lower Guinean Ichthyofaunal Province	478
Mann, David (see Schärer, Michelle T.)	
Marceñiuk, Alexandre P., Ricardo Betancur-R., Arturo Acero P., and Janice Muriel-Cunha. Review of the Genus <i>Cathorops</i> (Siluriformes: Ariidae) from the Caribbean and Atlantic South America, with Description of a New Species	77
Marinho, Manoela M. F. (see Martins, Fernanda O.)	
Maritz, Bryan, and Graham J. Alexander. Dwarfs on the Move: Spatial Ecology of the World's Smallest Viper, <i>Bitis schneideri</i>	115
Maritz, Bryan, and Graham J. Alexander. Movement Patterns in the Smallest Viper, <i>Bitis schneideri</i>	732
Marques, Otavio A. V. (see Bovo, Rafael P.)	
Martin, José (see Gabirot, Marianne)	
Martins, Fernanda O., Manoela M. F. Marinho, Francisco Langeani, and Jane P. Serra. A New Species of <i>Hypostomus</i> (Siluriformes: Loricariidae) from the Upper Rio Paraguay Basin, Brazil	494
Matamoros, Wilfredo A. Ecology and Evolution of Poeciliid Fishes. By Jonathan P. Evans, Andrea Pilastro, and Ingo Schlupp (eds.)	172
Mattingly, Hayden T. (see Bergen, Darrin R.)	
Maulany, Risma I., David T. Booth, and Greg S. Baxter. Emergence Success and Sex Ratio of Natural and Relocated Nests of Olive Ridley Turtles from Alas Purwo National Park, East Java, Indonesia	738
Meyer, Katherin P. (see Ennen, Joshua R.)	
Milko, Laura V. Integrating Museum and GIS Data to Identify Changes in Species Distributions Driven by a Disturbance-Induced Invasion	307
Miller, Gabriel J., Lora L. Smith, Steve A. Johnson, and Richard Franz. Home Range Size and Habitat Selection in the Florida Pine Snake (<i>Pituophis melanoleucus mugitus</i>)	706
Millsbaugh, Joshua J. (see Bodinof, Catherine M.)	
Mong, Tony (see Bodinof, Catherine M.)	
Montgomery, Chad E. (see Ray, Julie M.)	
Morgan, Margaret H. (see Duckett, Paul E.)	
Moriarty, John J. Turtles in Your Pocket: A Guide to Freshwater and Terrestrial Turtles of the Upper Midwest. By T. VanDeWalle. Frogs and Toads in Your Pocket: A Guide to Amphibians of the Upper Midwest. By T. VanDeWalle	357
Munroe, Thomas A. (see Duplain, René R.)	
Murdy, Edward O. (see Shibukawa, Koichi)	
Muriel-Cunha, Janice (see Marceñiuk, Alexandre P.)	
Mushet, David M., Ned H. Euliss, Jr., and Craig A. Stockwell. Mapping Anuran Habitat Suitability to Estimate Effects of Grassland and Wetland Conservation Programs	321
Nascimento, Juliana (see Provete, Diogo B.)	
Neely, David A. Peterson Field Guide to Freshwater Fishes: North America, North of Mexico. Second edition. By Lawrence M. Page and Brooks M. Burr	166
Nemeth, Michael I. (see Schärer, Michelle T.)	
Nickerson, Max A. (see Pitt, Amber L.)	
Nifong, James C., Adam E. Rosenblatt, Nathan A. Johnson, William Barichivich, Brian R. Silliman, and Michael R. Heithaus. American Alligator Digestion Rate of Blue Crabs and Its Implications for Stomach Contents Analysis	419
Nunes, Ivan, Axel Kwet, and José P. Pombal, Jr. Taxonomic Revision of the <i>Scinax alter</i> Species Complex (Anura: Hylidae)	554
O'Connor, Constance M., Erin E. Reardon, and Lauren J. Chapman. Shorter Gills in Mouth-brooding Females of the Cichlid <i>Pseudocrenilabrus multicolor</i>	382
O'Leary, Maureen A., Raphaël Sarr, Raymond Malou, El Hadji Sow, Chris Lepre, and Robert V. Hill. A New Fossil Amiid from the Eocene of Senegal and the Persistence of Extinct Marine Amiids after the Cretaceous-Paleogene Boundary	603
Oldfield, Ron. Sex Determination and Differentiation in Fish. By A. Cnaani (ed.)	776
Olson, Christina A., and Karen H. Beard. Diet of the Introduced Greenhouse Frog in Hawaii	121
Orr, James Wilder (see Pietsch, Theodore W.)	
Orr, James Wilder. Two New Species of Snailfishes of the Genus <i>Careproctus</i> (Scorpaeniformes: Liparidae) from the Bering Sea and Eastern North Pacific Ocean, with a Redescription of <i>Careproctus ovigerus</i>	257
Ory, Bridget A. (see Hughey, Myra C.)	
Osman, Amany M. (see Abdel-Aziz, Sayeda H.)	
Özen, Özcan (see Smith, David G.)	
Pabijan, Maciej (see Lehtinen, Richard M.)	
Pagnucco, Katie S., Cynthia A. Paszkowski, and Garry J. Scrimgeour. Characterizing Movement Patterns and	

- Spatio-temporal Use of Under-road Tunnels by Long-toed Salamanders in Waterton Lakes National Park, Canada 331
- Paszowski, Cynthia A. (see Pagnucco, Katie S.)
- Pechmann, Joseph H. K. (see Alexander, Laura G.)
- Pereira, Edson H. L. (see Reis, Roberto E.)
- Pereira, Edson H. L. (see Ribeiro, Alexandre C.)
- Pereira, Edson H. L., and Marcelo R. Britto. A New Distinctively Colored Catfish of the Genus *Pareiorhaphis* (Siluriformes: Loricariidae) from the Rio Piracicaba, Upper Rio Doce Basin, Brazil 519
- Peterman, William E., Grant M. Connette, Brett N. Spatola, Lori S. Eggert, and Raymond D. Semlitsch. Identification of Polymorphic Loci in *Ambystoma annulatum* and Review of Cross-species Microsatellite Use in the Genus *Ambystoma* 570
- Peterson, Mark S. (see Lang, Erik T.)
- Pianka, Eric R. Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles. By J. B. Losos 583
- Picerno, Pablo (see Gabirot, Marianne)
- Pietsch, Theodore W., James Wilder Orr, and William N. Eschmeyer. *Mustelus felis* Ayres, 1854, a Senior Synonym of the Leopard Shark, *Triakis semifasciata* Girard, 1855 (Carchariformes: Triakidae), Invalidated by "Reversal of Precedence" 98
- Pinto, Roberta R., and Ronaldo Fernandes. A New Blind Snake Species of the Genus *Tricheilostoma* from Espinhaço Range, Brazil and Taxonomic Status of *Rena dimidiata* (Jan, 1861) (Serpentes: Epictinae: Leptotyphlopidae) 37
- Pisani, George R., and Robert Powell. Joseph Collins (1939–2012) 351
- Pitt, Amber L., and Max A. Nickerson. Reassessment of the Turtle Community in the North Fork of White River, Ozark County, Missouri 367
- Pizzatto, Lígia. Reproductive Biology and Phylogeny of Snakes. By Robert D. Aldridge and David M. Sever (eds.) 355
- Polloni, Pamela (see Hartel, Karsten E.)
- Pombal, José P., Jr. (see Nunes, Ivan)
- Popescu, Viorel D., Bekka S. Brodie, Malcolm L. Hunter, and Joseph D. Zydlewski. Use of Olfactory Cues by Newly Metamorphosed Wood Frogs (*Lithobates sylvaticus*) during Emigration 424
- Powell, Robert (see Pisani, George R.)
- Pritchard, Peter C. H. Saving Sea Turtles: Extraordinary Stories from the Battle Against Extinction. By J. R. Spotila 588
- Provete, Diogo B., Michel V. Garey, Luís Felipe Toledo, Juliana Nascimento, Luciana B. Lourenço, Denise de C. Rossa-Feres, and Célio F. B. Haddad. Redescription of *Physalaemus barrooi* (Anura: Leiuperidae) 507
- Puglis, Holly J. (see Ramirez, Edward A.)
- Pyron, R. Alexander. Reptilian Cytogenetics and Genomics. By E. Olmo (ed.) 586
- Quagio-Grassiotto, Irani, Luiz R. Malabarba, Marco A. Azevedo, John R. Burns, Clarianna Martins Baicere-Silva, and Rodrigo Quevedo. Unique Derived Features in Spermiogenesis and Sperm Morphology Supporting a Close Relationship between the Species of *Hollandichthys* and *Rachoviscus* (Characiformes: Characidae) 609
- Qualls, Carl P. (see Selman, Will)
- Quevedo, Rodrigo (see Quagio-Grassiotto, Irani)
- Quiroga, Lorena B. (see Lobo, Fernando)
- Ramírez-Pinilla, Martha Patricia (see Valencia-Aguilar, Anyelet)
- Ramirez, Edward A., Holly J. Puglis, Alicia Ritzenthaler, and Michelle Boone. Terrestrial Movements and Habitat Preferences of Male Cricket Frogs on a Golf Course 191
- Rapp Py-Daniel, Lucia (see de Oliveira, Renildo R.)
- Rapp Py-Daniel, Lucia (see Rocha, Marcelo)
- Rasmussen, Arne R., Johan Elmberg, Kate L. Sanders, and Peter Gravlund. Rediscovery of the Rare Sea Snake *Hydrophis parviceps* Smith 1935: Identification and Conservation Status 276
- Ray, Julie M., Chad E. Montgomery, Heidi K. Mahon, Alan H. Savitzky, and Karen R. Lips. Goo-Eaters: Diets of the Neotropical Snakes *Dipsas* and *Sibon* in Central Panama 197
- Reardon, Erin E. (see O'Connor, Constance M.)
- Reis, Roberto E. (see Sant'Anna, Vivianne B.)
- Reis, Roberto E., Edson H. L. Pereira, and Pablo Lehmann A. A New Genus and Species of Hypoptopomatine Catfish (Siluriformes: Loricariidae) from the Upper Rio São Francisco Basin, Brazil 6
- Reques, Ricardo (see García-de-Lomas, Juan)
- Ribeiro, Alexandre C., Flávio C. T. Lima, and Edson H. L. Pereira. A New Genus and Species of a Minute Suckermouth Armored Catfish (Siluriformes: Loricariidae) from the Rio Tocantins Drainage, Central Brazil: The Smallest Known Loricariid Catfish 637
- Ribeiro, José W., Jr., Albertina P. Lima, and William E. Magnusson. The Effect of Riparian Zones on Species Diversity of Frogs in Amazonian Forests 375
- Ritzenthaler, Alicia (see Ramirez, Edward A.)
- Robertson, Jeanne M. Amphibians and Reptiles: An Introduction to Their Natural History and Conservation. By M. L. Crump 587
- Rocha, Carlos Frederico D. (see Sabagh, Leandro T.)
- Rocha, Luiz A. Groupers of the World: A Field and Market Guide. By Matthew T. Craig, Yvonne J. Sadovy de Mitcheson, and Phillip C. Heemstra 767
- Rocha, Marcelo S. (see de Oliveira, Renildo R.)
- Rocha, Marcelo, Henrique Lazzarotto, and Lucia Rapp Py-Daniel. A New Species of *Scoloplax* with a Remarkable New Tooth Morphology within Loricarioidea (Siluriformes: Scoloplacidae) 670
- Rodríguez-Robles, Javier A. (see Webber, Michael M.)
- Rodríguez, Mónica S., Marcel R. Cavallaro, and Matthew R. Thomas. A New Diminutive Species of *Loricaria* (Siluriformes: Loricariidae) from the Rio Paraguay System, Mato Grosso do Sul, Brazil 49
- Roosenburg, Willem M. (see Brown, Ashley D.)
- Rosales Casián, Jorge A. (see Baumann, Hannes)
- Rosenblatt, Adam E. (see Nifong, James C.)
- Rowell, Timothy J. (see Schäfer, Michelle T.)
- Ryan, Michael J. (see Dawson, Beth)
- Sabagh, Leandro T., Gustavo Luis Ferreira, Christina W. C. Branco, Carlos Frederico D. Rocha, and Natacha Y. N. Dias. Larval Diet in Bromeliad Pools: A Case Study of Tadpoles of Two Species in the Genus *Scinax* (Hylidae) 683
- Sanabria, Eduardo A. (see Lobo, Fernando)
- Sanders, Kate L. (see Rasmussen, Arne R.)
- Sant'Anna, Vivianne B., Maria Laura S. Delapieve, and Roberto E. Reis. A New Species of *Potamorhaphis* (Beloniformes: Belonidae) from the Amazon Basin 663

- Sarr, Raphaël (see O'Leary, Maureen A.)
- Savage, Jay M. Amphibians of Central America. By Gunther Köhler 170
- Savitzky, Alan H. (see Ray, Julie M.)
- Schall, Jos. J. Hemoparasites of the Reptilia: Color Atlas and Text. By S. R. Telford, Jr. 768
- Schärer, Michelle T., Michael I. Nemeth, David Mann, James Locascio, Richard S. Appeldoorn, and Timothy J. Rowell. Sound Production and Reproductive Behavior of Yellowfin Grouper, *Mycteroperca venenosa* (Serranidae) at a Spawning Aggregation 135
- Schmitter-Soto, Juan J. Historical Biogeography of Neotropical Freshwater Fishes. By James S. Albert and Roberto E. Reis (eds.) 162
- Schnell, Nalani K., and G. David Johnson. Ontogenetic Fusion of the Third and Fourth Pharyngobranchial in Barbeled Dragonfishes (Stomiidae, Teleostei) with a Revision of the Identity of the Single Posterior Upper Pharyngeal Toothplate 394
- Schoenfuss, Heiko L., and William F. Font. John Michael "Mike" Fitzsimons (1941–2012) 762
- Schuette, Chawna D. (see Bodinof, Catherine M.)
- Scrimgeour, Garry J. (see Pagnucco, Katie S.)
- Selman, Will, Jodie M. Jawor, and Carl P. Qualls. Seasonal Variation of Corticosterone Levels in *Graptomys flavimaculata*, an Imperiled Freshwater Turtle 698
- Semlitsch, Raymond D. (see Peterman, William E.)
- Senchal, Victoria (see Torres-Mejia, Mauricio)
- Serra, Jane P. (see Martins, Fernanda O.)
- Shaffer, H. Bradley. Biology of Turtles: From Structures to Strategies of Life. By Jeanette Wyneken, Matthew H. Godfrey, and Vincent Bels (eds.) 771
- Shea, Glenn M. The Eponym Dictionary of Reptiles. By B. Beolens, M. Watson, and M. Grayson 578
- Shibukawa, Koichi, and Edward O. Murdy. A Redescription of the Eel Goby *Trypauchenopsis* (Gobiidae: Amblyopinae) with Comments on Relationships 527
- Siegel, Dustin S. Reproduction of Amphibians. By M. Ogielska 168
- Silliman, Brian R. (see Nifong, James C.)
- Sinervo, Barry (see Corl, Ammon)
- Skelton, Paul H. (see Vari, Richard P.)
- Slack, William T. (see Lang, Erik T.)
- Smith, David G., Erhan Irmak, and Özcan Özen. A Redescription of the Eel *Panturichthys fowleri* (Anguilliformes: Heterenchelyidae), with a Synopsis of the Heterenchelyidae 484
- Smith, Lora L. (see Howze, Jennifer M.)
- Smith, Lora L. (see Miller, Gabriel J.)
- Sow, El Hadji (see O'Leary, Maureen A.)
- Sparks, John S., and David F. Gruber. A New Mesophotic Clingfish (Teleostei: Gobiessocidae) from the Bahamas 251
- Spatola, Brett N. (see Peterman, William E.)
- Stiassny, Melanie L. J. (see Mamonekene, Victor)
- Stockwell, Craig A. (see Mushet, David M.)
- Stow, Adam J. (see Duckett, Paul E.)
- Temple-Miller, Kathleen (see Brown, Ashley D.)
- Thomas, Matthew R. (see Rodriguez, Mónica S.)
- Toledo, Luis Felipe (see Provete, Diogo B.)
- Torres-Mejia, Mauricio, Emilton Hernández, and Victoria Senchal. A New Species of *Astyanacinus* (Characiformes: Characidae) from the Río Magdalena System, Colombia 501
- Torres, José María (see Garcia-de-Lomas, Juan)
- Uller, Tobias. Sex Determination and Differentiation in Reptiles. By Ettore Olmo (ed.) 773
- Valdecantos, María Soledad, Federico Arias, and Robert E. Espinoza. Herbivory in *Liolaemus poecilochromus*, a Small, Cold-Climate Lizard from the Andes of Argentina 203
- Valencia-Aguilar, Anyelet, Fernando Castro-Herrera, and Martha Patricia Ramírez-Pinilla. Microhabitats for Oviposition and Male Clutch Attendance in *Hyalinobatrachium aureoguttatum* (Anura: Centrolenidae) 722
- Valencia, Jorge (see Gabirot, Marianne)
- Vari, Richard P. (see de Santana, Carlos David)
- Vari, Richard P. (see Fernández, Luis)
- Vari, Richard P., Carl J. Ferraris, Jr., and Paul H. Skelton. New Species of *Congolani* (Siluriformes: Amphiliidae) from the Southern Congo River Basin 626
- Vences, Miguel (see Lehtinen, Richard M.)
- Walter, Ryan P. (see Lindstrom, Daniel P.)
- Wanner, Mark D. (see Bodinof, Catherine M.)
- Wassersug, Richard J. (see Calich, Hannah J.)
- Webber, Michael M., Xavier Glaudas, and Javier A. Rodríguez-Robles. Do Sidewinder Rattlesnakes (*Crotalus cerastes*, Viperidae) Cease Feeding During the Breeding Season? 100
- Weight, Lee A. (see Baldwin, Carole C.)
- White, Matthew M. (see Brown, Ashley D.)
- Wilkinson, Jeffery A. (see Dever, Jennifer A.)
- Yagi, Katharine T., and Jacqueline D. Litzgus. The Effects of Flooding on the Spatial Ecology of Spotted Turtles (*Clemmys guttata*) in a Partially Mined Peatland 179
- Yerman, Michelle N. (see Leis, Jeffrey M.)
- Zuanon, Jansen (see de Oliveira, Renildo R.)
- Zydlewski, Joseph D. (see Popescu, Viorel D.)